

SOCIETY OF ACTUARIES/CASUALTY ACTUARIAL SOCIETY

EXAM C CONSTRUCTION AND EVALUATION OF ACTUARIAL MODELS

EXAM C SAMPLE SOLUTIONS

Copyright 2008 by the Society of Actuaries and the Casualty Actuarial Society

Some of the questions in this study note are taken from past SOA/CAS examinations.

Question #1**Key: E**

The 40th percentile is the .4(12) = 4.8th smallest observation. By interpolation it is .2(86) + .8(90) = 89.2. The 80th percentile is the .8(12) = 9.6th smallest observation. By interpolation it is .4(200) + .6(210) = 206.

The equations to solve are

$$.4 = \frac{(89.2/\theta)^\gamma}{1+(89.2/\theta)^\gamma} \text{ and } .8 = \frac{(206/\theta)^\gamma}{1+(206/\theta)^\gamma} .$$

Solving each for the parenthetical expression gives $\frac{2}{3} = (89.2/\theta)^\gamma$ and $4 = (206/\theta)^\gamma$.

Taking the ratio of the second equation to the first gives $6 = (206/89.2)^\gamma$ which leads to $\gamma = \ln(6)/\ln(206/89.2) = 2.1407$. Then $4^{1/2.1407} = 206/\theta$ for $\theta = 107.8$.

Question #2**Key: E**

The standard for full credibility is $\left(\frac{1.645}{.02}\right)^2 \left(1 + \frac{\text{Var}(X)}{E(X)^2}\right)$ where X is the claim size

variable. For the Pareto variable, $E(X) = .5/5 = .1$ and $\text{Var}(X) = \frac{2(.5)^2}{5(4)} - (.1)^2 = .015$.

Then the standard is $\left(\frac{1.645}{.02}\right)^2 \left(1 + \frac{.015}{.1^2}\right) = 16,913$ claims.

Question #3**Key: B**

The kernel is a triangle with a base of 4 and a height at the middle of 0.5 (so the area is 1). The length of the base is twice the bandwidth. Any observation within 2 of 2.5 will contribute to the estimate. For the observation at 2, when the triangle is centered at 2, the height of the triangle at 2.5 is .375 (it is one-quarter the way from 2 to the end of the triangle at 4 and so the height is one-quarter the way from 0.5 to 0). Similarly the points at 3 are also 0.5 away and so the height of the associated triangle is also .375. Each triangle height is weighted by the empirical probability at the associated point. So the estimate at 2.5 is $(1/5)(3/8) + (3/5)(3/8) + (1/5)(0) = 12/40$.

Question #4**Key: A**

The distribution function is $F(x) = \int_1^x \alpha t^{-\alpha-1} dt = -t^{-\alpha} \Big|_1^x = 1 - x^{-\alpha}$. The likelihood function is

$$\begin{aligned} L &= f(3)f(6)f(14)[1 - F(25)]^2 \\ &= \alpha 3^{-\alpha-1} \alpha 6^{-\alpha-1} \alpha 14^{-\alpha-1} (25^{-\alpha})^2 \\ &\propto \alpha^3 [3(6)(14)(625)]^{-\alpha}. \end{aligned}$$

Taking logs, differentiating, setting equal to zero, and solving:

$$\ln L = 3 \ln \alpha - \alpha \ln 157,500 \text{ plus a constant}$$

$$(\ln L)' = 3\alpha^{-1} - \ln 157,500 = 0$$

$$\hat{\alpha} = 3 / \ln 157,500 = .2507.$$

Question #5**Key: C**

$$\pi(q | 1,1) \propto p(1 | q)p(1 | q)\pi(q) = 2q(1-q)2q(1-q)4q^3 \propto q^5(1-q)^2$$

$$\int_0^1 q^5(1-q)^2 dq = 1/168, \quad \pi(q | 1,1) = 168q^5(1-q)^2.$$

The expected number of claims in a year is $E(X | q) = 2q$ and so the Bayesian estimate is

$$E(2q | 1,1) = \int_0^1 2q(168)q^5(1-q)^2 dq = 4/3.$$

The answer can be obtained without integrals by recognizing that the posterior distribution of q is beta with $a = 6$ and $b = 3$. The posterior mean is

$$E(q | 1,1) = a/(a+b) = 6/9 = 2/3. \text{ The posterior mean of } 2q \text{ is then } 4/3.$$

Question #6**Key: D**

For the method of moments estimate,

$$386 = e^{\mu + .5\sigma^2}, \quad 457,480.2 = e^{2\mu + 2\sigma^2}$$

$$5.9558 = \mu + .5\sigma^2, \quad 13.0335 = 2\mu + 2\sigma^2$$

$$\hat{\mu} = 5.3949, \quad \hat{\sigma}^2 = 1.1218.$$

Then

$$\begin{aligned}
E(X \wedge 500) &= e^{5.3949+5(1.1218)} \Phi\left(\frac{\ln 500 - 5.3949 - 1.1218}{\sqrt{1.1218}}\right) + 500 \left[1 - \Phi\left(\frac{\ln 500 - 5.3949}{\sqrt{1.1218}}\right)\right] \\
&= 386\Phi(-.2853) + 500[1 - \Phi(.7739)] \\
&= 386(.3877) + 500(.2195) = 259.
\end{aligned}$$

Note-these calculations use exact normal probabilities. Rounding and using the normal table that accompanies the exam will produce a different numerical answer but the same letter answer.

Question #7
DELETED

Question #8
Key: C

Let N be the Poisson claim count variable, let X be the claim size variable, and let S be the aggregate loss variable.

$$\mu(\theta) = E(S | \theta) = E(N | \theta)E(X | \theta) = \theta 10\theta = 10\theta^2$$

$$v(\theta) = \text{Var}(S | \theta) = E(N | \theta)E(X^2 | \theta) = \theta 200\theta^2 = 200\theta^3$$

$$\mu = E(10\theta^2) = \int_1^\infty 10\theta^2 (5\theta^{-6}) d\theta = 50/3$$

$$EPV = E(200\theta^3) = \int_1^\infty 200\theta^3 (5\theta^{-6}) d\theta = 500$$

$$VHM = \text{Var}(10\theta^2) = \int_1^\infty (10\theta^2)^2 (5\theta^{-6}) d\theta - (50/3)^2 = 222.22$$

$$k = 500 / 222.22 = 2.25.$$

Question #9
DELETED

Question #10
DELETED

Question #11**Key: D**

$$\Pr(\theta = 1 | X = 5) = \frac{f(5 | \theta = 1) \Pr(\theta = 1)}{f(5 | \theta = 1) \Pr(\theta = 1) + f(5 | \theta = 3) \Pr(\theta = 3)}$$

$$= \frac{(1/36)(1/2)}{(1/36)(1/2) + (3/64)(1/2)} = 16/43$$

$$\Pr(X_2 > 8 | X_1 = 5) = \Pr(X_2 > 8 | \theta = 1) \Pr(\theta = 1 | X_1 = 5) + \Pr(X_2 > 8 | \theta = 3) \Pr(\theta = 3 | X_1 = 5)$$

$$= (1/9)(16/43) + (3/11)(27/43) = .2126.$$

For the last line, $\Pr(X > 8 | \theta) = \int_8^{\infty} \theta(x + \theta)^{-2} dx = \theta(8 + \theta)^{-1}$ is used.

Question #12**Key: C**

The sample mean for X is 720 and for Y is 670. The mean of all 8 observations is 695.

$$(730 - 720)^2 + (800 - 720)^2 + (650 - 720)^2 + (700 - 720)^2$$

$$+ (655 - 670)^2 + (650 - 670)^2 + (625 - 670)^2 + (750 - 670)^2$$

$$\hat{v} = \frac{3475}{2(4 - 1)} = 3475$$

$$\hat{a} = \frac{(720 - 695)^2 + (670 - 695)^2}{2 - 1} - \frac{3475}{4} = 381.25$$

$$\hat{k} = 3475 / 381.25 = 9.1148$$

$$\hat{Z} = \frac{4}{4 + 9.1148} = .305$$

$$P_c = .305(670) + .695(695) = 687.4.$$

Question #13**Key: B**

There are 430 observations. The expected counts are $430(.2744) = 117.99$, $430(.3512) = 151.02$, $430(.3744) = 160.99$. The test statistic is

$$\frac{(112 - 117.99)^2}{117.99} + \frac{(180 - 151.02)^2}{151.02} + \frac{(138 - 160.99)^2}{160.99} = 9.15.$$

Question #14**Key: B**

From the information, the asymptotic variance of $\hat{\theta}$ is $1/4n$. Then

$$\text{Var}(2\hat{\theta}) = 4\text{Var}(\hat{\theta}) = 4(1/4n) = 1/n.$$

Note that the delta method is not needed for this problem, although using it leads to the same answer.

Question #15

Key: A

The posterior probability density is

$$\pi(p | 1,1,1,1,1,1,1,1) \propto \Pr(1,1,1,1,1,1,1,1 | p)\pi(p) \propto p^8(2) \propto p^8.$$

$$\pi(p | 1,1,1,1,1,1,1,1) = \frac{p^8}{\int_0^5 p^8 dp} = \frac{p^8}{(.5^9)/9} = 9(.5^{-9})p^8.$$

$$\begin{aligned} \Pr(X_9 = 1 | 1,1,1,1,1,1,1,1) &= \int_0^5 \Pr(X_9 = 1 | p)\pi(p | 1,1,1,1,1,1,1,1)dp \\ &= \int_0^5 p9(.5^{-9})p^8 dp = 9(.5^{-9})(.5^{10})/10 = .45. \end{aligned}$$

Question #16

Key: A

$${}_3\hat{p}_1 = \frac{18}{27} \frac{26}{32} \frac{20}{25} = \frac{13}{30}. \text{ Greenwood's approximation is}$$

$$\left(\frac{13}{30}\right)^2 \left(\frac{9}{18(27)} + \frac{6}{26(32)} + \frac{5}{20(25)}\right) = .0067.$$

Question #17**Key: D**

$$\hat{H}(3) = 5/30 + 9/27 + 6/32 = 0.6875$$

$$\hat{V}ar(\hat{H}(3)) = 5/(30)^2 + 9/(27)^2 + 6/(32)^2 = 0.02376$$

The 95% log-transformed confidence interval is:

$$\hat{H}(3)U, \text{ where } U = \exp\left(\pm \frac{1.96\sqrt{.02376}}{.6875}\right) = \exp(\pm 0.43945)$$

The confidence interval is:

$$[0.6875 \exp(-0.43945), 0.6875 \exp(0.43945)] = [0.443, 1.067].$$

Question #18**Key: D**

The means are $.5(250) + .3(2,500) + .2(60,000) = 12,875$ and $.7(250) + .2(2,500) + .1(60,000) = 6,675$ for risks 1 and 2 respectively.

The variances are $.5(250)^2 + .3(2,500)^2 + .2(60,000)^2 - 12,875^2 = 556,140,625$ and $.7(250)^2 + .2(2,500)^2 + .1(60,000)^2 - 6,675^2 = 316,738,125$ respectively.

The overall mean is $(2/3)(12,875) + (1/3)(6,675) = 10,808.33$ and so

$EPV = (2/3)(556,140,625) + (1/3)(316,738,125) = 476,339,792$ and

$VHM = (2/3)(12,875)^2 + (1/3)(6,675)^2 - 10,808.33^2 = 8,542,222$. Then,

$k = 476,339,792/8,542,222 = 55.763$ and $Z = 1/(1 + 55.763) = .017617$.

The credibility estimate is $.017617(250) + .982383(10,808.33) = 10,622$.

Question #19**Key: D**

The first two sample moments are 15 and 500, and the first two population moments are

$E(X) = .5(\theta + \sigma)$ and $E(X^2) = .5(2\theta^2 + 2\sigma^2) = \theta^2 + \sigma^2$. These can be obtained either

through integration or by recognizing the density function as a two-point mixture of

exponential densities. The equations to solve are $30 = \theta + \sigma$ and $500 = \theta^2 + \sigma^2$. From

the first equation, $\sigma = 30 - \theta$ and substituting into the second equation gives

$500 = \theta^2 + (30 - \theta)^2 = 2\theta^2 - 60\theta + 900$. The quadratic equation has two solutions, 10 and

20. Because $\theta > \sigma$ the answer is 20.

Question #20**Key: D**

There are four possible samples, (5,5), (5,9), (9,5), and (9,9). For each, the estimator g must be calculated. The values are 0, 4, 4, and 0 respectively. Assuming a population in which the values 5 and 9 each occur with probability .5, the population variance is $.5(5-7)^2 + .5(9-7)^2 = 4$. The mean square error is approximated as $.25[(0-4)^2 + (4-4)^2 + (4-4)^2 + (0-4)^2] = 8$.

Question #21**Key: B**

From the Poisson distribution, $\mu(\lambda) = \lambda$ and $v(\lambda) = \lambda$. Then,

$\mu = E(\lambda) = 6/100 = .06$, $EPV = E(\lambda) = .06$, $VHM = Var(\lambda) = 6/100^2 = .0006$ where the various moments are evaluated from the gamma distribution. Then, $k = .06/.0006 = 100$ and $Z = 450/(450 + 100) = 9/11$ where the 450 is the total number of insureds contributing experience. The credibility estimate of the expected number of claims for one insured in month 4 is $(9/11)(25/450) + (2/11)(.06) = .056364$. For 300 insureds the expected number of claims is $300(.056364) = 16.9$.

Question #22**Key: C**

The likelihood function is $L(\alpha, \theta) = \prod_{j=1}^{200} \frac{\alpha \theta^\alpha}{(x_j + \theta)^{\alpha+1}}$ and its logarithm is

$l(\alpha, \theta) = 200 \ln(\alpha) + 200\alpha \ln(\theta) - (\alpha + 1) \sum_{i=1}^{200} \ln(x_i + \theta)$. When evaluated at the hypothesized values of 1.5 and 7.8, the loglikelihood is -821.77 . The test statistic is $2(821.77 - 817.92) = 7.7$. With two degrees of freedom (0 free parameters in the null hypothesis versus 2 in the alternative), the test statistic falls between the 97.5th percentile (7.38) and the 99th percentile (9.21).

Question #23**Key: E**

Assume that $\theta > 5$. Then the expected counts for the three intervals are $15(2/\theta) = 30/\theta$, $15(3/\theta) = 45/\theta$, and $15(\theta - 5)/\theta = 15 - 75/\theta$ respectively. The quantity to minimize is

$$\frac{1}{5}[(30\theta^{-1} - 5)^2 + (45\theta^{-1} - 5)^2 + (15 - 75\theta^{-1} - 5)^2].$$

Differentiating (and ignoring the coefficient of 1/5) gives the equation

$$-2(30\theta^{-1} - 5)30\theta^{-2} - 2(45\theta^{-1} - 5)45\theta^{-2} + 2(10 - 75\theta^{-1})75\theta^{-2} = 0.$$

Multiplying through by θ^3 and dividing by 2 reduces the equation to

$$-(30 - 5\theta)30 - (45 - 5\theta)45 + (10\theta - 75)75 = -8550 + 1125\theta = 0$$

$$\text{for a solution of } \hat{\theta} = 8550/1125 = 7.6.$$

Question #24**Key: E**

$\pi(\theta | 1) \propto \theta(1.5\theta^5) \propto \theta^{1.5}$. The required constant is the reciprocal of $\int_0^1 \theta^{1.5} d\theta = \theta^{2.5} / 2.5 \Big|_0^1 = .4$

and so $\pi(\theta | 1) = 2.5\theta^{1.5}$. The requested probability is

$$\Pr(\theta > .6 | 1) = \int_{.6}^1 2.5\theta^{1.5} d\theta = \theta^{2.5} \Big|_{.6}^1 = 1 - .6^{2.5} = .721.$$

Question #25**Key: A**

k	kn_k / n_{k-1}
0	
1	0.81
2	0.92
3	1.75
4	2.29
5	2.50
6	3.00

Positive slope implies that the negative binomial distribution is a good choice. Alternatively, the sample mean and variance are 1.2262 and 1.9131 respectively. With the variance substantially exceeding the mean, the negative binomial model is again supported.

Question #26**Key: B**

The likelihood function is $\frac{e^{-1/(2\theta)}}{2\theta} \cdot \frac{e^{-2/(2\theta)}}{2\theta} \cdot \frac{e^{-3/(2\theta)}}{2\theta} \cdot \frac{e^{-15/(3\theta)}}{3\theta} = \frac{e^{-8/\theta}}{24\theta^4}$. The loglikelihood function is $-\ln 24 - 4\ln(\theta) - 8/\theta$. Differentiating with respect to θ and setting the result equal to 0 yields $-\frac{4}{\theta} + \frac{8}{\theta^2} = 0$ which produces $\hat{\theta} = 2$.

Question #27**Key: E**

The absolute difference of the credibility estimate from its expected value is to be less than or equal to $k\mu$ (with probability P). That is,

$$\left| [ZX_{\text{partial}} + (1-Z)M] - [Z\mu + (1-Z)M] \right| \leq k\mu$$

$$-k\mu \leq ZX_{\text{partial}} - Z\mu \leq k\mu.$$

Adding μ to all three sides produces answer choice (E).

Question #28**Key: C**

In general,

$$E(X^2) - E[(X \wedge 150)^2] = \int_0^{200} x^2 f(x) dx - \int_0^{150} x^2 f(x) dx - 150^2 \int_{150}^{200} f(x) dx = \int_{150}^{200} (x^2 - 150^2) f(x) dx.$$

Assuming a uniform distribution, the density function over the interval from 100 to 200 is $6/7400$ (the probability of $6/74$ assigned to the interval divided by the width of the interval). The answer is

$$\int_{150}^{200} (x^2 - 150^2) \frac{6}{7400} dx = \left(\frac{x^3}{3} - 150^2 x \right) \frac{6}{7400} \Big|_{150}^{200} = 337.84.$$

Question #29**Key: B**

The probabilities are from a binomial distribution with 6 trials. Three successes were observed.

$$\Pr(3 | \text{I}) = \binom{6}{3} (.1)^3 (.9)^3 = .01458, \Pr(3 | \text{II}) = \binom{6}{3} (.2)^3 (.8)^3 = .08192,$$

$$\Pr(3 | \text{III}) = \binom{6}{3} (.4)^3 (.6)^3 = .27648$$

The probability of observing three successes is $.7(.01458) + .2(.08192) + .1(.27648) = .054238$. The three posterior probabilities are:

$$\Pr(\text{I} | 3) = \frac{.7(.01458)}{.054238} = .18817, \Pr(\text{II} | 3) = \frac{.2(.08192)}{.054238} = .30208, \Pr(\text{III} | 3) = \frac{.1(.27648)}{.054238} = .50975.$$

The posterior probability of a claim is then

$$.1(.18817) + .2(.30208) + .4(.50975) = .28313.$$

Question #30

Key: E

$.542 = \hat{F}(n) = 1 - e^{-\hat{H}(n)}$, $\hat{H}(n) = .78$. The Nelson-Aalen estimate is the sum of successive s/r values. From the problem statement, $r = 100$ at all surrender times while the s -values follow the pattern 1, 2, 3, Then,

$$.78 = \frac{1}{100} + \frac{2}{100} + \dots + \frac{n}{100} = \frac{n(n+1)}{200} \text{ and the solution is } n = 12.$$

Question # 31

Answer: C

$$g = [12(.45)] = [5.4] = 5; \quad h = 5.4 - 5 = 0.4.$$

$$\hat{\pi}_{.45} = .6x_{(5)} + .4x_{(6)} = .6(360) + .4(420) = 384.$$

Question # 32

Answer: D

N is distributed *Poisson*(λ)

$$\mu = E(\lambda) = \alpha\theta = 1(1.2) = 1.2.$$

$$v = E(\lambda) = 1.2; \quad a = \text{Var}(\lambda) = \alpha\theta^2 = 1(1.2)^2 = 1.44.$$

$$k = \frac{1.2}{1.44} = \frac{5}{6}; \quad Z = \frac{2}{2 + 5/6} = \frac{12}{17}.$$

Thus, the estimate for Year 3 is

$$\frac{12}{17}(1.5) + \frac{5}{17}(1.2) = 1.41.$$

Note that a Bayesian approach produces the same answer.

Question # 33**Answer: C**

At the time of the second failure,

$$\hat{H}(t) = \frac{1}{n} + \frac{1}{n-1} = \frac{23}{132} \Rightarrow n = 12.$$

At the time of the fourth failure,

$$\hat{H}(t) = \frac{1}{12} + \frac{1}{11} + \frac{1}{10} + \frac{1}{9} = .3854.$$

Question # 34**Answer: B**

The likelihood is:

$$L = \prod_{j=1}^n \frac{r(r+1)\cdots(r+x_j-1)\beta^{x_j}}{x_j!(1+\beta)^{r+x_j}} \propto \prod_{j=1}^n \beta^{x_j} (1+\beta)^{-r-x_j}.$$

The loglikelihood is:

$$l = \sum_{j=1}^n \left[x_j \ln \beta - (r+x_j) \ln(1+\beta) \right]$$

$$l' = \sum_{j=1}^n \left[\frac{x_j}{\beta} - \frac{r+x_j}{1+\beta} \right] = 0$$

$$0 = \sum_{j=1}^n \left[x_j(1+\beta) - (r+x_j)\beta \right] = \sum_{j=1}^n x_j - rn\beta$$

$$0 = n\bar{x} - rn\beta; \quad \hat{\beta} = \bar{x}/r.$$

Question # 35**Answer: C**

The Bühlmann credibility estimate is $Zx + (1-Z)\mu$ where x is the first observation. The Bühlmann estimate is the least squares approximation to the Bayesian estimate. Therefore, Z and μ must be selected to minimize

$$\frac{1}{3}[Z + (1-Z)\mu - 1.5]^2 + \frac{1}{3}[2Z + (1-Z)\mu - 1.5]^2 + \frac{1}{3}[3Z + (1-Z)\mu - 3]^2.$$

Setting partial derivatives equal to zero will give the values. However, it should be clear that μ is the average of the Bayesian estimates, that is,

$$\mu = \frac{1}{3}(1.5 + 1.5 + 3) = 2.$$

The derivative with respect to Z is (deleting the coefficients of 1/3):

$$2(-Z + .5)(-1) + 2(.5)(0) + 2(Z - 1)(1) = 0$$

$$Z = .75.$$

The answer is

$$.75(1) + .25(2) = 1.25.$$

Question # 36

Answer: E

The confidence interval is $(\hat{S}(t_0)^{1/\theta}, \hat{S}(t_0)^\theta)$.

Taking logarithms of both endpoints gives the two equations

$$\ln .695 = -.36384 = \frac{1}{\theta} \ln \hat{S}(t_0)$$

$$\ln .843 = -.17079 = \theta \ln \hat{S}(t_0).$$

Multiplying the two equations gives

$$.06214 = [\ln \hat{S}(t_0)]^2$$

$$\ln \hat{S}(t_0) = -.24928$$

$$\hat{S}(t_0) = .77936.$$

The negative square root is required in order to make the answer fall in the interval (0,1).

Question # 37**Answer: B**

The likelihood is:

$$L = \frac{\alpha 150^\alpha}{(150 + 225)^{\alpha+1}} \frac{\alpha 150^\alpha}{(150 + 525)^{\alpha+1}} \frac{\alpha 150^\alpha}{(150 + 950)^{\alpha+1}}$$

$$= \frac{\alpha^3 150^{3\alpha}}{(375 \cdot 675 \cdot 1100)^{\alpha+1}}.$$

The loglikelihood is:

$$l = 3 \ln \alpha + 3\alpha \ln 150 - (\alpha + 1) \ln(375 \cdot 675 \cdot 1100)$$

$$l' = \frac{3}{\alpha} + 3 \ln 150 - \ln(375 \cdot 675 \cdot 1100) = \frac{3}{\alpha} - 4.4128$$

$$\hat{\alpha} = 3 / 4.4128 = .6798.$$

Question # 38**Answer: D**For this problem, $r = 4$ and $n = 7$. Then,

$$\hat{v} = \frac{33.60}{4(7-1)} = 1.4 \text{ and } \hat{a} = \frac{3.3}{4-1} - \frac{1.4}{7} = .9.$$

Then,

$$k = \frac{1.4}{.9} = \frac{14}{9}; \quad Z = \frac{7}{7 + (14/9)} = \frac{63}{77} = .82.$$

Question # 39**Answer: B** X is the random sum $Y_1 + Y_2 + \dots + Y_N$. N has a negative binomial distribution with $r = \alpha = 1.5$ and $\beta = \theta = 0.2$.

$$E(N) = r\beta = 0.3$$

$$\text{Var}(N) = r\beta(1 + \beta) = 0.36$$

$$E(Y) = 5000$$

$$\text{Var}(Y) = 25,000,000$$

$$E(X) = 0.3 \times 5000 = 1500$$

$$Var(X) = 0.3 \times 25,000,000 + 0.36 \times 25,000,000 = 16,500,000$$

Number of exposures (insureds) required for full credibility

$$n_{FULL} = (1.645 / 0.05)^2 \times 16,500,000 / (1500)^2 = 7937.67.$$

Number of expected claims required for full credibility

$$E(N) \times n_{FULL} = 0.3 \times 7937.67 = 2381.$$

Question # 40

Answer: E

X	$F_n(x)$	$F_n(x^-)$	$F_0(x)$	$ F_n(x) - F_0(x) $	$ F_n(x^-) - F_0(x) $
29	0.2	0	0.252	0.052	0.252
64	0.4	0.2	0.473	0.073	0.273
90	0.6	0.4	0.593	0.007	0.193
135	0.8	0.6	0.741	0.059	0.141
182	1.00	0.8	0.838	0.162	0.038

where:

$$\hat{\theta} = \bar{x} = 100 \text{ and } F_0(x) = 1 - e^{-x/100}.$$

The maximum value from the last two columns is 0.273.

Question # 41

Answer: E

$$\mu = E(\lambda) = 1; \quad v = E(\sigma^2) = 1.25; \quad a = Var(\lambda) = 1/12.$$

$$k = v/a = 15; \quad Z = \frac{1}{1+15} = \frac{1}{16}.$$

Thus, the estimate for Year 2 is

$$\frac{1}{16}(0) + \frac{15}{16}(1) = .9375.$$

Question # 42
DELETED

Question # 43
Answer: E

The posterior density, given an observation of 3 is:

$$\begin{aligned}\pi(\theta | 3) &= \frac{f(3 | \theta)\pi(\theta)}{\int_1^{\infty} f(3 | \theta)\pi(\theta)d\theta} = \frac{\frac{2\theta^2}{(3+\theta)^3} \frac{1}{\theta^2}}{\int_1^{\infty} 2(3+\theta)^{-3} d\theta} \\ &= \frac{2(3+\theta)^{-3}}{-(3+\theta)^{-2} \Big|_1^{\infty}} = 32(3+\theta)^{-3}, \quad \theta > 1.\end{aligned}$$

Then,

$$\Pr(\Theta > 2) = \int_2^{\infty} 32(3+\theta)^{-3} d\theta = -16(3+\theta)^{-2} \Big|_2^{\infty} = \frac{16}{25} = .64.$$

Question # 44
Answer: B

$$\begin{aligned}L &= F(1000)^7 [F(2000) - F(1000)]^6 [1 - F(2000)]^7 \\ &= (1 - e^{-1000/\theta})^7 (e^{-1000/\theta} - e^{-2000/\theta})^6 (e^{-2000/\theta})^7 \\ &= (1 - p)^7 (p - p^2)^6 (p^2)^7 \\ &= p^{20} (1 - p)^{13}\end{aligned}$$

where $p = e^{-1000/\theta}$. The maximum occurs at $p = 20/33$ and so $\hat{\theta} = -1000 / \ln(20/33) = 1996.90$.

Question # 45
Answer: A

$$E(X | \theta) = \theta / 2.$$

$$\begin{aligned}E(X_3 | 400, 600) &= \int_{600}^{\infty} E(X | \theta) f(\theta | 400, 600) d\theta = \int_{600}^{\infty} \frac{\theta}{2} 3 \frac{600^3}{\theta^4} d\theta = \frac{3(600^3)}{2} \frac{\theta^{-2}}{-2} \Big|_{600}^{\infty} \\ &= \frac{3(600^3)(600^{-2})}{4} = 450.\end{aligned}$$

Question # 46**Answer: D**

The data may be organized as follows:

T	Y	d	$\hat{S}(t)$
2	10	1	$(9/10) = .9$
3	9	2	$.9(7/9) = .7$
5	7	1	$.7(6/7) = .6$
6	5	1	$.6(4/5) = .48$
7	4	1	$.48(3/4) = .36$
9	2	1	$.36(1/2) = .18$

Because the product-limit estimate is constant between observations, the value of $\hat{S}(8)$ is found from $\hat{S}(7) = .36$.

Question # 47**Answer: C**

The maximum likelihood estimate for the Poisson distribution is the sample mean:

$$\hat{\lambda} = \bar{x} = \frac{50(0) + 122(1) + 101(2) + 92(3)}{365} = 1.6438.$$

The table for the chi-square test is:

Number of days	Probability	Expected*	Chi-square
0	$e^{-1.6438} = .19324$	70.53	5.98
1	$1.6438e^{-1.6438} = .31765$	115.94	0.32
2	$\frac{1.6438^2 e^{-1.6438}}{2} = .26108$	95.30	0.34
3+	.22803**	83.23	0.92

*365x(Probability) **obtained by subtracting the other probabilities from 1

The sum of the last column is the test statistic of 7.56. Using 2 degrees of freedom (4 rows less 1 estimated parameter less 1) the model is rejected at the 2.5% significance level but not at the 1% significance level.

Question # 48**Answer: D**

$$\mu(0) = \frac{.4(0) + .1(1) + .1(2)}{.6} = .5; \quad \mu(1) = \frac{.1(0) + .2(1) + .1(2)}{.4} = 1$$

$$\mu = .6(.5) + .4(1) = .7$$

$$a = .6(.5^2) + .4(1^2) - .7^2 = .06$$

$$v(0) = \frac{.4(0) + .1(1) + .1(4)}{.6} - .5^2 = \frac{7}{12}; \quad v(1) = \frac{.1(0) + .2(1) + .1(4)}{.4} - 1^2 = .5$$

$$v = .6(7/12) + .4(.5) = 11/20$$

$$k = v/a = 55/6; \quad Z = \frac{10}{10 + 55/6} = \frac{60}{115}$$

$$\text{Bühlmann credibility premium} = \frac{60}{115} \frac{10}{10} + \frac{55}{115} (.7) = .8565.$$

Question # 49**Answer: C**

$$\mu = .5(0) + .3(1) + .1(2) + .1(3) = .8$$

$$\sigma^2 = .5(0) + .3(1) + .1(4) + .1(9) - .64 = .96$$

$$E(S_n^2) = \frac{n-1}{n} \sigma^2 = \frac{3}{4} (.96) = .72$$

$$\text{bias} = .72 - .96 = -.24.$$

Question # 50**Answer: C**

The four classes have means .1, .2, .5, and .9 respectively and variances .09, .16, .25, and .09 respectively.

Then,

$$\mu = .25(.1 + .2 + .5 + .9) = .425$$

$$v = .25(.09 + .16 + .25 + .09) = .1475$$

$$a = .25(.01 + .04 + .25 + .81) - .425^2 = .096875$$

$$k = .1475 / .096875 = 1.52258$$

$$Z = \frac{4}{4 + 1.52258} = .7243$$

The estimate is $[.7243(2/4) + .2757(.425)] \cdot 5 = 2.40$.

Question # 51
DELETED

Question # 52
Answer: A

The distribution used for simulation is given by the observed values.

Question # 53
Answer: B

First obtain the distribution of aggregate losses:

Value	Probability
0	1/5
25	(3/5)(1/3) = 1/5
100	(1/5)(2/3)(2/3) = 4/45
150	(3/5)(2/3) = 2/5
250	(1/5)(2)(2/3)(1/3) = 4/45
400	(1/5)(1/3)(1/3) = 1/45

$$\mu = (1/5)(0) + (1/5)(25) + (4/45)(100) + (2/5)(150) + (4/45)(250) + (1/45)(400) = 105$$

$$\begin{aligned} \sigma^2 &= (1/5)(0^2) + (1/5)(25^2) + (4/45)(100^2) + (2/5)(150^2) \\ &\quad + (4/45)(250^2) + (1/45)(400^2) - 105^2 = 8,100. \end{aligned}$$

Question # 54
Answer: A

Loss Range	Cum. Prob.
0 – 100	0.320
100 – 200	0.530
200 – 400	0.800

400 – 750	0.960
750 – 1000	0.980
1000 – 1500	1.000

At 400, $F(x) = 0.8 = 1 - e^{-\frac{400}{\theta}}$; solving gives $\theta = 248.53$.

Question # 55

Answer: B

$$\Pr(\text{class1} | 1) = \frac{(1/2)(1/3)}{(1/2)(1/3) + (1/3)(1/6) + (1/6)(0)} = \frac{3}{4}$$

$$\Pr(\text{class2} | 1) = \frac{(1/3)(1/6)}{(1/2)(1/3) + (1/3)(1/6) + (1/6)(0)} = \frac{1}{4}$$

$$\Pr(\text{class3} | 1) = \frac{(1/6)(0)}{(1/2)(1/3) + (1/3)(1/6) + (1/6)(0)} = 0$$

because the prior probabilities for the three classes are 1/2, 1/3, and 1/6 respectively.

The class means are

$$\mu(1) = (1/3)(0) + (1/3)(1) + (1/3)(2) = 1$$

$$\mu(2) = (1/6)(1) + (2/3)(2) + (1/6)(3) = 2.$$

The expectation is

$$E(X_2 | 1) = (3/4)(1) + (1/4)(2) = 1.25.$$

Question # 56

Answer: E

The first, second, third, and sixth payments were observed at their actual value and each contributes $f(x)$ to the likelihood function. The fourth and fifth payments were paid at the policy limit and each contributes $1 - F(x)$ to the likelihood function. This is answer (E).

Question #57

Answer is E

For an interval running from c to d , the uniform density function is $f(x) = g/[n(d-c)]$ where g is the number of observations in the interval and n is the sample size. The contribution to the second raw moment for this interval is:

$$\int_c^d x^2 \frac{g}{n(d-c)} dx = \frac{gx^3}{3n(d-c)} \Big|_c^d = \frac{g(d^3 - c^3)}{3n(d-c)}.$$

For this problem, the second raw moment is:

$$\frac{1}{90} \left[\frac{30(25^3 - 0^3)}{3(25 - 0)} + \frac{32(50^3 - 25^3)}{3(50 - 25)} + \frac{20(100^3 - 50^3)}{3(100 - 50)} + \frac{8(200^3 - 100^3)}{3(200 - 100)} \right] = 3958.33.$$

Question #58

Answer is B

Because the Bayes and Bühlmann results must be identical, this problem can be solved either way. For the Bühlmann approach, $\mu(\lambda) = v(\lambda) = \lambda$. Then, noting that the prior distribution is a gamma distribution with parameters 50 and 1/500, we have:

$$\mu = E(\lambda) = 50/500 = 0.1$$

$$v = E(\lambda) = 0.1$$

$$a = \text{Var}(\lambda) = 50/500^2 = 0.0002$$

$$k = v/a = 500$$

$$Z = 1500/(1500 + 500) = 0.75$$

$$\bar{X} = \frac{75 + 210}{600 + 900} = 0.19.$$

The credibility estimate is $0.75(0.19) + 0.25(0.1) = 0.1675$. For 1100 policies, the expected number of claims is $1100(0.1675) = 184.25$.

For the Bayes approach, the posterior density is proportional to (because in a given year the number of claims has a Poisson distribution with parameter λ times the number of policies)

$$\frac{e^{-600\lambda} (600\lambda)^{75}}{75!} \frac{e^{-900\lambda} (900\lambda)^{210}}{210!} \frac{(500\lambda)^{50} e^{-500\lambda}}{\lambda \Gamma(50)} \propto \lambda^{335} e^{-2000\lambda}$$

which is a gamma density with parameters 335 and 1/2000. The expected number of claims per policy is $335/2000 = 0.1675$ and the expected number of claims in the next year is 184.25.

Question #59

Answer is E

The q - q plot takes the ordered values and plots the j th point at $j/(n+1)$ on the horizontal axis and at $F(x_j; \theta)$ on the vertical axis. For small values, the model assigns more probability to being below that value than occurred in the sample. This indicates that the model has a heavier left tail than the data. For large values, the model again assigns more probability to being below that value (and so less probability to being above that value). This indicates that the model has a lighter right tail than the data. Of the five answer choices, only E is consistent with these observations. In addition, note that as you go from 0.4 to 0.6 on the horizontal axis (thus looking at the middle 20% of the data), the q - q plot increases from about 0.3 to 0.4 indicating that the model puts only about 10% of the probability in this range, thus confirming answer E.

Question #60

Answer is C

The posterior probability of having one of the coins with a 50% probability of heads is proportional to $(.5)(.5)(.5)(.5)(4/6) = 0.04167$. This is obtained by multiplying the probabilities of making the successive observations 1, 1, 0, and 1 with the 50% coin times the prior probability of 4/6 of selecting this coin. The posterior probability for the 25% coin is proportional to $(.25)(.25)(.75)(.25)(1/6) = 0.00195$ and the posterior probability for the 75% coin is proportional to $(.75)(.75)(.25)(.75)(1/6) = 0.01758$. These three numbers total 0.06120. Dividing by this sum gives the actual posterior probabilities of 0.68088, 0.03186, and 0.28726. The expected value for the fifth toss is then $(.68088)(.5) + (.03186)(.25) + (.28726)(.75) = 0.56385$.

Question #61

Answer is A

Because the exponential distribution is memoryless, the excess over the deductible is also exponential with the same parameter. So subtracting 100 from each observation yields data from an exponential distribution and noting that the maximum likelihood estimate is the sample mean gives the answer of 73.

Working from first principles,

$$L(\theta) = \frac{f(x_1)f(x_2)f(x_3)f(x_4)f(x_5)}{[1 - F(100)]^5} = \frac{\theta^{-1}e^{-125/\theta}\theta^{-1}e^{-150/\theta}\theta^{-1}e^{-165/\theta}\theta^{-1}e^{-175/\theta}\theta^{-1}e^{-250/\theta}}{(e^{-100/\theta})^5}$$

$$= \theta^{-5}e^{-365/\theta}.$$

Taking logarithms and then a derivative gives

$$l(\theta) = -5 \ln(\theta) - 365/\theta, l'(\theta) = -5/\theta + 365/\theta^2 = 0.$$

The solution is $\hat{\theta} = 365/5 = 73$.

Question #62**Answer is D**

The number of claims for each insured has a binomial distribution with $n = 1$ and q unknown. We have

$$\mu(q) = q, v(q) = q(1 - q)$$

$$\mu = E(q) = 0.1, \text{ given in item (iv)}$$

$$a = \text{Var}(q) = E(q^2) - E(q)^2 = E(q^2) - 0.01 = 0.01, \text{ given in item (v)}$$

$$\text{Therefore, } E(q^2) = 0.02$$

$$v = E(q - q^2) = 0.1 - 0.02 = 0.08$$

$$k = v/a = 8, Z = \frac{10}{10+8} = 5/9.$$

Then the expected number of claims in the next one year is $(5/9)(0) + (4/9)(0.1) = 2/45$ and the expected number of claims in the next five years is $5(2/45) = 2/9 = 0.22$.

Question #63**DELETED****Question #64****Answer is E**

The model distribution is $f(x|\theta) = 1/\theta, 0 < x < \theta$. Then the posterior distribution is proportional to

$$\pi(\theta | 400, 600) \propto \frac{1}{\theta} \frac{1}{\theta} \frac{500}{\theta^2} \propto \theta^{-4}, \theta > 600.$$

It is important to note the range. Being a product, the posterior density function is non-zero only when all three terms are non-zero. Because one of the observations was equal to 600, the value of the parameter must be greater than 600 in order for the density function at 600 to be positive. Or, by general reasoning, posterior probability can only be assigned to possible values. Having observed the value 600 we know that parameter values less than or equal to 600 are not possible.

The constant is obtained from $\int_{600}^{\infty} \theta^{-4} d\theta = \frac{1}{3(600)^3}$ and thus the exact posterior density is

$\pi(\theta | 400, 600) = 3(600)^3 \theta^{-4}, \theta > 600$. The posterior probability of an observation exceeding 550 is

$$\begin{aligned}\Pr(X_3 > 550 | 400, 600) &= \int_{600}^{\infty} \Pr(X_3 > 550 | \theta) \pi(\theta | 400, 600) d\theta \\ &= \int_{600}^{\infty} \frac{\theta - 550}{\theta} 3(600)^3 \theta^{-4} d\theta = 0.3125\end{aligned}$$

where the first term in the integrand is the probability of exceeding 550 from the uniform distribution.

Question #65

Answer is C

$$E(N) = r\beta = 0.40$$

$$\text{Var}(N) = r\beta(1 + \beta) = 0.48$$

$$E(Y) = \theta / (\alpha - 1) = 500$$

$$\text{Var}(Y) = \theta^2 \alpha / [(\alpha - 1)^2 (\alpha - 2)] = 750,000$$

Therefore,

$$E(X) = 0.40(500) = 200$$

$$\text{Var}(X) = 0.40(750,000) + 0.48(500)^2 = 420,000$$

The full credibility standard is $n = \left(\frac{1.645}{0.05}\right)^2 \frac{420,000}{200^2} = 11,365$ and then

$$Z = \sqrt{2500/11,365} = 0.47.$$

Question #66

Answer is E

The sample variance is $s^2 = \frac{(1-3)^2 + (2-3)^2 + (3-3)^2 + (4-3)^2 + (5-3)^2}{4} = 2.5$. The estimator of $E[X]$ is the sample mean and the variance of the sample mean is the variance divided by the sample size, estimated here as $2.5/n$. Setting the standard deviation of the estimator equal to 0.05 gives the equation $\sqrt{2.5/n} = 0.05$ which yields $n = 1000$.

Question #67

Answer is E

$$\mu(r) = E(X | r) = E(N)E(Y) = r\beta\theta / (\alpha - 1) = 100r$$

$$\begin{aligned}v(r) &= \text{Var}(X | r) = \text{Var}(N)E(Y)^2 + E(N)\text{Var}(Y) \\ &= r\beta(1 + \beta)\theta^2 / (\alpha - 1)^2 + r\beta\alpha\theta^2 / [(\alpha - 1)^2 (\alpha - 2)] = 210,000r.\end{aligned}$$

$$v = E(210,000r) = 210,000(2) = 420,000$$

$$a = \text{Var}(100r) = (100)^2(4) = 40,000$$

$$k = v/a = 10.5$$

$$Z = 100/(100+10.5) = 0.905.$$

Question #68

Answer is B

$$\text{Using all participants, } S^T(4) = \left(1 - \frac{35}{300}\right) \left(1 - \frac{74}{265}\right) \left(1 - \frac{34}{191}\right) \left(1 - \frac{32}{157}\right) = 0.41667.$$

$$\text{Using only Country B, } S^B(4) = \left(1 - \frac{15}{100}\right) \left(1 - \frac{20}{85}\right) \left(1 - \frac{20}{65}\right) \left(1 - \frac{10}{45}\right) = 0.35.$$

$$\text{The difference is, } S^T(4) - S^B(4) = 0.41667 - 0.35 = 0.0667 = 0.07.$$

Question #69

Answer is B

For an exponential distribution the maximum likelihood estimate of the mean is the sample mean. We have

$$E(\bar{X}) = E(X) = \theta, \text{Var}(\bar{X}) = \text{Var}(X)/n = \theta^2/n.$$

$$cv = SD(\bar{X})/E(\bar{X}) = [\theta/\sqrt{n}]/\theta = 1/\sqrt{n} = 1/\sqrt{5} = 0.447.$$

If the above facts are not known, the loglikelihood function can be used:

$$L(\theta) = \theta^{-n} e^{-\sum x_j/\theta}, \quad l(\theta) = -n \ln \theta - n\bar{X}/\theta, \quad l'(\theta) = -n\theta^{-1} + n\bar{X}\theta^{-2} = 0 \Rightarrow \hat{\theta} = \bar{X}.$$

$$l''(\theta) = n\theta^{-2} - 2n\bar{X}\theta^{-3}, \quad I(\theta) = E[-n\theta^{-2} + 2n\bar{X}\theta^{-3}] = n\theta^{-2}.$$

$$\text{Then, } \text{Var}(\hat{\theta}) = \theta^2/n.$$

Question #70

Answer is D

Because the total expected claims for business use is 1.8, it must be that 20% of business users are rural and 80% are urban. Thus the unconditional probabilities of being business-rural and business-urban are 0.1 and 0.4 respectively. Similarly the probabilities of being pleasure-rural and pleasure-urban are also 0.1 and 0.4 respectively. Then,

$$\mu = 0.1(1.0) + 0.4(2.0) + 0.1(1.5) + 0.4(2.5) = 2.05$$

$$v = 0.1(0.5) + 0.4(1.0) + 0.1(0.8) + 0.4(1.0) = 0.93$$

$$a = 0.1(1.0^2) + 0.4(2.0^2) + 0.1(1.5^2) + 0.4(2.5^2) - 2.05^2 = 0.2225$$

$$k = v/a = 4.18$$

$$Z = 1/(1 + 4.18) = 0.193.$$

Question #71

Answer is A

No. claims	Hypothesize d	Observe d	Chi-square
1	250	235	$15^2/250 = 0.90$
2	350	335	$15^2/350 = 0.64$
3	240	250	$10^2/240 = 0.42$
4	110	111	$1^2/110 = 0.01$
5	40	47	$7^2/40 = 1.23$
6+	10	22	$12^2/10 = 14.40$

The last column sums to the test statistic of 17.60 with 5 degrees of freedom (there were no estimated parameters), so from the table reject at the 0.005 significance level.

Question #72

Answer is C

In part (ii) you are given that $\mu = 20$. In part (iii) you are given that $a = 40$. In part (iv) you are given that $v = 8,000$. Therefore, $k = v/a = 200$. Then,

$$\bar{X} = \frac{800(15) + 600(10) + 400(5)}{1800} = \frac{100}{9}$$

$$Z = \frac{1800}{1800 + 200} = 0.9$$

$$P_c = 0.9(100/9) + 0.1(20) = 12.$$

Question #73**Answer is C**

$$\Pr(X > 30,000) = S(30,000) = \left(1 - \frac{1}{10 - 2/2}\right) \left(1 - \frac{1}{7 - 2/2}\right) = 20/27 = 0.741.$$

Question #74**DELETED****Question #75****Answer is D**

$$E(X) = \int_{\delta}^{\infty} \frac{x}{\theta} e^{-(x-\delta)/\theta} dx = \int_0^{\infty} \frac{y+\delta}{\theta} e^{-y/\theta} dy = \theta + \delta$$

$$E(X^2) = \int_{\delta}^{\infty} \frac{x^2}{\theta} e^{-(x-\delta)/\theta} dx = \int_0^{\infty} \frac{y^2 + 2y\delta + \delta^2}{\theta} e^{-y/\theta} dy = 2\theta^2 + 2\theta\delta + \delta^2.$$

Both derivations use the substitution $y = x - \delta$ and then recognize that the various integrals are requesting moments from an ordinary exponential distribution. The method of moments solves the two equations

$$\theta + \delta = 10$$

$$2\theta^2 + 2\theta\delta + \delta^2 = 130.6$$

producing $\hat{\delta} = 4.468$.

It is faster to do the problem if it is noted that $X = Y + \delta$ where Y has an ordinary exponential distribution. Then $E(X) = E(Y) + \delta = \theta + \delta$ and $\text{Var}(X) = \text{Var}(Y) = \theta^2$.

Question #76**Answer is D**

The posterior density is proportional to the product of the probability of the observed value and the prior density. Thus, $\pi(\theta | N > 0) \propto \Pr(N > 0 | \theta)\pi(\theta) = (1 - e^{-\theta})\theta e^{-\theta}$.

The constant of proportionality is obtained from $\int_0^{\infty} \theta e^{-\theta} - \theta e^{-2\theta} d\theta = \frac{1}{1^2} - \frac{1}{2^2} = 0.75$.

The posterior density is $\pi(\theta | N > 0) = (4/3)(\theta e^{-\theta} - \theta e^{-2\theta})$.

Then,

$$\begin{aligned}\Pr(N_2 > 0 | N_1 > 0) &= \int_0^\infty \Pr(N_2 > 0 | \theta) \pi(\theta | N_1 > 0) d\theta = \int_0^\infty (1 - e^{-\theta})(4/3)(\theta e^{-\theta} - \theta e^{-2\theta}) d\theta \\ &= \frac{4}{3} \int_0^\infty \theta e^{-\theta} - 2\theta e^{-2\theta} + \theta e^{-3\theta} d\theta = \frac{4}{3} \left(\frac{1}{1^2} - \frac{2}{2^2} + \frac{1}{3^2} \right) = 0.8148.\end{aligned}$$

Question #77

Answer is E

The interval is centered at 2.09 and the plus/minus term is 0.46 which must equal $1.96\hat{\sigma}$ and so $\hat{\sigma} = 0.2347$. For the log-transformed interval we need

$\phi = e^{1.96(0.2347)/2.09} = 1.2462$. The lower limit is $2.09/1.2462 = 1.68$ and the upper limit is $2.09(1.2462) = 2.60$.

Question #78

Answer is B

From item (ii), $\mu = 1000$ and $a = 50$. From item (i), $v = 500$. Therefore, $k = v/a = 10$ and $Z = 3/(3+10) = 3/13$. Also, $\bar{X} = (750 + 1075 + 2000)/3 = 1275$. Then

$$P_c = (3/13)(1275) + (10/13)(1000) = 1063.46.$$

Question #79

Answer is C

$$f(x) = p \frac{1}{100} e^{-x/100} + (1-p) \frac{1}{10,000} e^{-x/10,000}$$

$$L(100, 200) = f(100)f(2000)$$

$$= \left(\frac{pe^{-1}}{100} + \frac{(1-p)e^{-0.01}}{10,000} \right) \left(\frac{pe^{-20}}{100} + \frac{(1-p)e^{-0.2}}{10,000} \right)$$

Question #80

DELETED

Question # 81**Answer: C**

Which distribution is it from?

0.25 < 0.30, so it is from the exponential.

Given that Y is from the exponential, we want

$$\Pr(Y \leq y) = F(y) = 0.69$$

$$1 - e^{-y/\theta} = 0.69$$

$$1 - e^{-y/0.5} = 0.69 \text{ since mean} = 0.5$$

$$\frac{-y}{0.5} = \ln(1 - 0.69) = -1.171$$

$$y = 0.5855$$

Question #82**Key: B**

If you happen to remember this distribution from the Simulation text (example 4d in third edition), you could use:

$$n = \text{Int} \left(\frac{\log(1-u)}{\log q} \right) + 1 = \text{Int} \frac{\log 0.95}{\log 0.1} + 1 = 0 + 1 = 1$$

For mere mortals, you get the simulated value of N from the definition of the inverse transformation method:

$$f(1) = F(1) = 0.9$$

$$0.05 \leq 0.9 \text{ so } n = 1$$

$$x_1 = \frac{1}{\lambda} \log^{(1-v_1)} = -\frac{1}{0.01} \log 0.7 = 35.67$$

The amount of total claims during the year = 35.67

Question #83**Key: B**

$$F(0) = 0.8$$

$$F(t) = 0.8 + 0.00005(t-1000), \quad 1000 \leq t \leq 5000$$

$$0.75 \Rightarrow 0 \text{ found since } F(0) \geq 0.75$$

0.85 \Rightarrow 2000 found since $F(2000) = 0.85$
Average of those two outcomes is 1000.

Question #84

Key: A

$$B = \begin{cases} c(400 - x) & x < 400 \\ 0 & x \geq 400 \end{cases}$$

$$100 = E(B) = c \cdot 400 - cE(X \wedge 400)$$

$$= c \cdot 400 - c \cdot 300 \left(1 - \frac{300}{300 + 400} \right)$$

$$= c \left(400 - 300 \cdot \frac{4}{7} \right)$$

$$c = \frac{100}{228.6} = 0.44$$

Question #85

Key: C

Let $N = \#$ of computers in department

Let $X =$ cost of a maintenance call

Let $S =$ aggregate cost

$$\text{Var}(X) = [\text{Standard Deviation}(X)]^2 = 200^2 = 40,000$$

$$E(X^2) = \text{Var}(X) + [E(X)]^2 \\ = 40,000 + 80^2 = 46,400$$

$$E(S) = N \times \lambda \times E(X) = N \times 3 \times 80 = 240N$$

$$\text{Var}(S) = N \times \lambda \times E(X^2) = N \times 3 \times 46,400 = 139,200N$$

We want $0.1 \geq \Pr(S > 1.2E(S))$

$$\geq \Pr\left(\frac{S - E(S)}{\sqrt{139,200N}} > \frac{0.2E(S)}{\sqrt{139,200N}}\right) \Rightarrow \frac{0.2 \times 240N}{373.1\sqrt{N}} \geq 1.282 = \Phi(0.9)$$

$$N \geq \left(\frac{1.282 \times 373.1}{48}\right)^2 = 99.3$$

Question #86

Key: D

The modified severity, X^* , represents the conditional payment amount given that a payment occurs. Given that a payment is required ($X > d$), the payment must be uniformly distributed between 0 and $c \cdot (b - d)$.

The modified frequency, N^* , represents the number of losses that result in a payment.

The deductible eliminates payments for losses below d , so only $1 - F_x(d) = \frac{b-d}{b}$ of losses will require payments. Therefore, the Poisson parameter for the modified frequency distribution is $\lambda \cdot \frac{b-d}{b}$. (Reimbursing $c\%$ after the deductible affects only the payment amount and not the frequency of payments).

Question #87

Key: E

$$f(x) = 0.01, \quad 0 \leq x \leq 80$$

$$= 0.01 - 0.00025(x - 80) = 0.03 - 0.00025x, \quad 80 < x \leq 120$$

$$E(x) = \int_0^{80} 0.01x \, dx + \int_{80}^{120} (0.03x - 0.00025x^2) \, dx$$

$$= \frac{0.01x^2}{2} \Big|_0^{80} + \frac{0.03x^2}{2} \Big|_{80}^{120} - \frac{0.00025x^3}{3} \Big|_{80}^{120}$$

$$= 32 + 120 - 101.33 = 50.66667$$

$$E(X - 20)_+ = E(X) - \int_0^{20} x f(x) \, dx - 20(1 - \int_0^{20} f(x) \, dx)$$

$$= 50.6667 - \frac{0.01x^2}{2} \Big|_0^{20} - 20(1 - 0.01x \Big|_0^{20})$$

$$= 50.6667 - 2 - 20(0.8) = 32.6667$$

$$\text{Loss Elimination Ratio} = 1 - \frac{32.6667}{50.6667} = 0.3553$$

Question #88**Key: B**

First restate the table to be CAC's cost, after the 10% payment by the auto owner:

Towing Cost, x	$p(x)$
72	50%
90	40%
144	10%

$$\text{Then } E(X) = 0.5 * 72 + 0.4 * 90 + 0.1 * 144 = 86.4$$

$$E(X^2) = 0.5 * 72^2 + 0.4 * 90^2 + 0.1 * 144^2 = 7905.6$$

$$\text{Var}(X) = 7905.6 - 86.4^2 = 440.64$$

$$\text{Because Poisson, } E(N) = \text{Var}(N) = 1000$$

$$E(S) = E(X)E(N) = 86.4 * 1000 = 86,400$$

$$\text{Var}(S) = E(N)\text{Var}(X) + E(X)^2\text{Var}(N) = 1000 * 440.64 + 86.4^2 * 1000 = 7,905,600$$

$$\Pr(S > 90,000) + \Pr\left(\frac{S - E(S)}{\sqrt{\text{Var}(S)}} > \frac{90,000 - 86,400}{\sqrt{7,905,600}}\right) = \Pr(Z > 1.28) = 1 - \Phi(1.28) = 0.10$$

Since the frequency is Poisson, you could also have used

$$\text{Var}(S) = \lambda E(X^2) = (1000)(7905.6) = 7,905,600$$

That way, you would not need to have calculated $\text{Var}(X)$.

Question #89**Key: C**

$$\text{LER} = \frac{E(X \wedge d)}{E(X)} = \frac{\theta(1 - e^{-d/\theta})}{\theta} = 1 - e^{-d/\theta}$$

$$\text{Last year } 0.70 = 1 - e^{-d/\theta} \Rightarrow -d = \theta \log 0.30$$

$$\text{Next year: } -d_{\text{new}} = \theta \log(1 - \text{LER}_{\text{new}})$$

$$\text{Hence } \theta \log(1 - \text{LER}_{\text{new}}) = -d_{\text{new}} = \frac{4}{3} \theta \log 0.30$$

$$\log(1 - \text{LER}_{\text{new}}) = -1.6053$$

$$(1 - \text{LER}_{\text{new}}) = e^{-1.6053} = 0.20$$

$$\text{LER}_{\text{new}} = 0.80$$

Question # 90**Answer: E**

The distribution of claims (a gamma mixture of Poissons) is negative binomial.

$$E(N) = E_{\Lambda}(E(N|\Lambda)) = E_{\Lambda}(\Lambda) = 3$$

$$Var(N) = E_{\Lambda}(Var(N|\Lambda)) + Var_{\Lambda}(E(N|\Lambda))$$

$$= E_{\Lambda}(\Lambda) + Var_{\Lambda}(\Lambda) = 6$$

$$r\beta = 3$$

$$r\beta(1 + \beta) = 6$$

$$(1 + \beta) = 6/3 = 2; \quad \beta = 1$$

$$r\beta = 3$$

$$r = 3$$

$$p_0 = (1 + \beta)^{-r} = 0.125$$

$$p_1 = \frac{r\beta}{(1 + \beta)^{r+1}} = 0.1875$$

$$\begin{aligned} \text{Pr ob(at most 1)} &= p_0 + p_1 \\ &= 0.3125 \end{aligned}$$

Question # 91**Answer: A**

$$E(S) = E(N) \times E(X) = 110 \times 1,101 = 121,110$$

$$Var(S) = E(N) \times Var(X) + E(X)^2 \times Var(N)$$

$$= 110 \times 70^2 + 1101^2 \times 750$$

$$= 909,689,750$$

$$\text{Std Dev } (S) = 30,161$$

$$\begin{aligned}\Pr(S < 100,000) &= \Pr(Z < (100,000 - 121,110) / 30,161) \text{ where } Z \text{ has standard normal} \\ &\text{distribution} \\ &= \Pr(Z < -0.70) = 0.242\end{aligned}$$

Question # 92
Answer: C

Let N = number of prescriptions then $S = N \times 40$

n	$f_N(n)$	$F_N(n)$	$1 - F_N(n)$
0	0.2000	0.2000	0.8000
1	0.1600	0.3600	0.6400
2	0.1280	0.4880	0.5120
3	0.1024	0.5904	0.4096

$$E(N) = 4 = \sum_{j=0}^{\infty} (1 - F(j))$$

$$\begin{aligned}E[(S - 80)_+] &= 40 \times E[(N - 2)_+] = 40 \times \sum_{j=2}^{\infty} (1 - F(j)) \\ &= 40 \times \left[\sum_{j=0}^{\infty} (1 - F(j)) - \sum_{j=0}^1 (1 - F(j)) \right] \\ &= 40(4 - 1.44) = 40 \times 2.56 = 102.40\end{aligned}$$

$$\begin{aligned}E[(S - 120)_+] &= 40 \times E[(N - 3)_+] = 40 \times \sum_{j=3}^{\infty} (1 - F(j)) \\ &= 40 \times \left[\sum_{j=0}^{\infty} (1 - F(j)) - \sum_{j=0}^2 (1 - F(j)) \right] \\ &= 40(4 - 1.952) = 40 \times 2.048 = 81.92\end{aligned}$$

Since no values of S between 80 and 120 are possible,

$$E[(S - 100)_+] = \frac{(120 - 100) \times E[(S - 80)_+] + (100 - 80) \times E[(S - 120)_+]}{120} = 92.16$$

Alternatively,

$$E[(S - 100)_+] = \sum_{j=0}^{\infty} (40j - 100)f_N(j) + 100f_N(0) + 60f_N(1) + 20f_N(2)$$

(The correction terms are needed because $(40j - 100)$ would be negative for $j = 0, 1, 2$; we need to add back the amount those terms would be negative)

$$\begin{aligned}
 &= 40 \sum_{j=0}^{\infty} j \times f_N(j) - 100 \sum_{j=0}^{\infty} f_N(j) + (100)(0.200) + (0.16)(60) + (0.128)(20) \\
 &= 40 E(N) - 100 + 20 + 9.6 + 2.56 \\
 &= 160 - 67.84 = 92.16
 \end{aligned}$$

Question #93

Answer: E

Method 1:

In each round,

N = result of first roll, to see how many dice you will roll

X = result of for one of the N dice you roll

S = sum of X for the N dice

$$E(X) = E(N) = 3.5$$

$$Var(X) = Var(N) = 2.9167$$

$$E(S) = E(N) * E(X) = 12.25$$

$$Var(S) = E(N)Var(X) + Var(N)E(X)^2$$

$$= (3.5)(2.9167) + (2.9167)(3.5)^2$$

$$= 45.938$$

Let S_{1000} = the sum of the winnings after 1000 rounds

$$E(S_{1000}) = 1000 * 12.25 = 12,250$$

$$Stddev(S_{1000}) = \text{sqrt}(1000 * 45.938) = 214.33$$

After 1000 rounds, you have your initial 15,000, less payments of 12,500, plus winnings of S_{1000} .

Since actual possible outcomes are discrete, the solution tests for continuous outcomes greater than 15000-0.5. In this problem, that continuity correction has negligible impact.

$$\Pr(15000 - 12500 + S_{1000} > 14999.5) =$$

$$= \Pr\left(\frac{S_{1000} - 12250}{214.33} > \frac{(14999.5 - 2500 - 12250)}{214.33}\right) =$$

$$= 1 - \Phi(1.17) = 0.12$$

Method 2

Realize that you are going to determine N 1000 times and roll the sum of those 1000 N 's dice, adding the numbers showing.

Let N_{1000} = sum of those N 's

$$E(N_{1000}) = 1000E(N) = (1000)(3.5) = 3500$$

$$\text{Var}(N_{1000}) = 1000\text{Var}(N) = 2916.7$$

$$E(S_{1000}) = E(N_{1000})E(X) = (3500)(3.5) = 12.250$$

$$\text{Var}(S_{1000}) = E(N_{1000})\text{Var}(X) + \text{Var}(N_{1000})E(X)^2$$

$$= (3500)(2.9167) + (2916.7)(3.5)^2 = 45.938$$

$$\text{Stddev}(S_{1000}) = 214.33$$

Now that you have the mean and standard deviation of S_{1000} (same values as method 1), use the normal approximation as shown with method 1.

Question #94

Answer: B

$$p_k = \left(a + \frac{b}{k}\right)p_{k-1}$$

$$0.25 = (a + b) \times 0.25 \Rightarrow a + b = 1$$

$$0.1875 = \left(a + \frac{b}{2}\right) \times 0.25 \Rightarrow \left(1 - \frac{b}{2}\right) \times 0.25 = 0.1875$$

$$b = 0.5$$

$$a = 0.5$$

$$p_3 = \left(0.5 + \frac{0.5}{3}\right) \times 0.1875 = 0.125$$

Question #95

Answer: E

$$\beta = \text{mean} = 4; \quad p_k = \beta^k / (1 + \beta)^{k+1}$$

n	$P(N = n)$
0	0.2
1	0.16
2	0.128
3	0.1024

x	$f^{(1)}(x)$	$f^{(2)}(x)$	$f^{(3)}(x)$
0	0	0	0
1	0.25	0	0
2	0.25	0.0625	0
3	0.25	0.125	0.0156

$f^{(k)}(x)$ = probability that, given exactly k claims occur, that the aggregate amount is x .

$f^{(1)}(x) = f(x)$; the claim amount distribution for a single claim

$$f^{(k)}(x) = \sum_{j=0}^x \left(f^{(k-1)}(j) \right) x f(x-j)$$

$$f_s(x) = \sum_{k=0}^x P(N = k) \times f^{(k)}(x); \text{ upper limit of sum is really } \infty, \text{ but here with smallest}$$

possible claim size 1, $f^{(k)}(x) = 0$ for $k > x$

$$f_s(0) = 0.2$$

$$f_s(1) = 0.16 * 0.25 = 0.04$$

$$f_s(2) = 0.16 * 0.25 + 0.128 * 0.0625 = 0.048$$

$$f_s(3) = 0.16 * 0.25 + 0.128 * 0.125 + 0.1024 * 0.0156 = 0.0576$$

$$F_s(3) = 0.2 + 0.04 + 0.048 + 0.0576 = 0.346$$

Question #96**Answer: E**Let L = incurred losses; P = earned premium = 800,000

$$\text{Bonus} = 0.15 \times \left(0.60 - \frac{L}{P} \right) \times P \text{ if positive}$$

$$= 0.15 \times (0.60P - L) \text{ if positive}$$

$$= 0.15 \times (480,000 - L) \text{ if positive}$$

$$= 0.15 \times (480,000 - (L \wedge 480,000))$$

$$E(\text{Bonus}) = 0.15 (480,000 - E(L \wedge 480,000))$$

From Appendix A.2.3.1

$$= 0.15 \{ 480,000 - [500,000 \times (1 - (500,000 / (480,000 + 500,000)))] \}$$

$$= 35,265$$

Question # 97**Key: D**

Model Solution:

Severity after increase	Severity after increase and deductible
60	0
120	20
180	80
300	200

$$\begin{aligned} \text{Expected payment per loss} &= 0.25 \times 0 + 0.25 \times 20 + 0.25 \times 80 + 0.25 \times 200 \\ &= 75 \end{aligned}$$

$$\begin{aligned} \text{Expected payments} &= \text{Expected number of losses} \times \text{Expected payment per loss} \\ &= 75 \times 300 \\ &= 22,500 \end{aligned}$$

Question # 98**Key: A**

Model Solution:

$$E(S) = E(N) E(X) = 50 \times 200 = 10,000$$

$$\begin{aligned}\text{Var}(S) &= E(N)\text{Var}(X) + E(X)^2\text{Var}(N) \\ &= (50)(400) + (200^2)(100) \\ &= 4,020,000\end{aligned}$$

$$\begin{aligned}\Pr(S < 8,000) &= \Pr\left(Z < \frac{8,000 - 10,000}{\sqrt{4,020,000}}\right) \\ &= \Pr(Z < -0.998) \cong 16\%\end{aligned}$$

Question #99

Key: B

Model Solution:

Let S denote aggregate losses before deductible.

$$E[S] = 2 \times 2 = 4, \text{ since mean severity is 2.}$$

$$f_S(0) = \frac{e^{-2}2^0}{0!} = 0.1353, \text{ since must have 0 number to get aggregate losses} = 0.$$

$$f_S(1) = \left(\frac{e^{-2}2}{1!}\right)\left(\frac{1}{3}\right) = 0.0902, \text{ since must have 1 loss whose size is 1 to get aggregate losses} = 1.$$

$$\begin{aligned}E(S \wedge 2) &= 0 \times f_S(0) + 1 \times f_S(1) + 2 \times (1 - f_S(0) - f_S(1)) \\ &= 0 \times 0.1353 + 1 \times 0.0902 + 2 \times (1 - 0.1353 - 0.0902) \\ &= 1.6392\end{aligned}$$

$$\begin{aligned}E[(S - 2)_+] &= E[S] - E[S \wedge 2] \\ &= 4 - 1.6392 \\ &= 2.3608\end{aligned}$$

Question #100

Key: C

Model Solution:

Limited expected value =

$$\int_0^{1000} (1 - F(x))dx = \int_0^{1000} (0.8e^{-0.02x} + 0.2e^{-0.001x})dx = \left(-40e^{-0.02x} - 200e^{-0.001x}\right)\Big|_0^{1000} = 40 + 126.4$$

$$= 166.4$$

Question #101

Key: B

Model Solution:

$$\begin{aligned}\text{Mean excess loss} &= \frac{E(X) - E(X \wedge 100)}{1 - F(100)} \\ &= \frac{331 - 91}{0.8} = 300\end{aligned}$$

$$E(X) = E(X \wedge 1000) \text{ since } F(1000) = 1.0$$

Question #102

Key: E

Model Solution:

$$\begin{aligned}\text{Expected insurance benefits per factory} &= E[(X - 1)_+] \\ &= 0.2 \times 1 + 0.1 \times 2 = 0.4.\end{aligned}$$

$$\text{Insurance premium} = (1.1) (2 \text{ factories}) (0.4 \text{ per factory}) = 0.88.$$

Let R = retained major repair costs, then

$$f_R(0) = 0.4^2 = 0.16$$

$$f_R(1) = 2 \times 0.4 \times 0.6 = 0.48$$

$$f_R(2) = 0.6^2 = 0.36$$

$$\begin{aligned}\text{Dividend} &= 3 - 0.88 - R - (0.15)(3), \text{ if positive} \\ &= 1.67 - R, \text{ if positive}\end{aligned}$$

$$E(\text{Dividend}) = (0.16)(1.67 - 0) + (0.48)(1.67 - 1) + (0.36)(0) = 0.5888$$

[The $(0.36)(0)$ term in the last line represents that with probability 0.36, $(1.67 - R)$ is negative so the dividend is 0.]

Question #103**Key: A**

Model Solution:

$$E[X] = \frac{\alpha\theta}{\alpha-1} = \frac{4\alpha}{\alpha-1} = 8 \Rightarrow 4\alpha = 8\alpha - 8$$

$$\alpha = 2$$

$$F(6) = 1 - \left(\frac{\theta}{6}\right)^\alpha = 1 - \left(\frac{4}{6}\right)^2$$

$$= 0.555$$

$$s(6) = 1 - F(6) = 0.444$$

Test Question: 104**Key: C**

$$E[N] = E_\Lambda[E[N|\Lambda]] = E_\Lambda[\Lambda] = 2$$

$$Var[N] = E_\Lambda[Var[N|\Lambda]] + Var_\Lambda[E[N|\Lambda]]$$

$$= E_\Lambda[\Lambda] + Var_\Lambda[\Lambda] = 2 + 2 = 4$$

Distribution is negative binomial (Loss Models, 3.3.2)

Per supplied tables

$$mean = r\beta = 2$$

$$Var = r\beta(1 + \beta) = 4$$

$$(1 + \beta) = 2$$

$$\beta = 1$$

$$r\beta = 2$$

$$r = 2$$

From tables

$$p_3 = \frac{r(r+1)(r+2)\beta^3}{3!(1+\beta)^{r+3}} = \frac{(2)(3)(4)1^3}{3!2^5} = \frac{4}{32} = 0.125$$

$$1000 p_3 = 125$$

Test Question: 105**Key: A**

Using the conditional mean and variance formulas:

$$E[N] = E_{\Lambda}(N|\Lambda)$$

$$Var[N] = Var_{\Lambda}(E(N|\Lambda)) + E_{\Lambda}(Var(N|\Lambda))$$

Since N , given Λ , is just a Poisson distribution, this simplifies to:

$$E[N] = E_{\Lambda}(\Lambda)$$

$$Var[N] = Var_{\Lambda}(\Lambda) + E_{\Lambda}(\Lambda)$$

We are given that $E[N] = 0.2$ and $Var[N] = 0.4$, subtraction gives $Var(\Lambda) = 0.2$ **Test Question: 106****Key: B** N = number of salmon X = eggs from one salmon S = total eggs.

$$E(N) = 100t$$

$$Var(N) = 900t$$

$$E(S) = E(N)E(X) = 500t$$

$$Var(S) = E(N)Var(X) + E^2(X)Var(N) = 100t \cdot 5 + 25 \cdot 900t = 23,000t$$

$$P(S > 10,000) = P\left(\frac{S - 500t}{\sqrt{23,000t}} > \frac{10,000 - 500t}{\sqrt{23,000t}}\right) = .95 \Rightarrow$$

$$10,000 - 500t = -1.645 \cdot \sqrt{23,000} \sqrt{t} = -250\sqrt{t}$$

$$40 - 2t = -\sqrt{t}$$

$$2(\sqrt{t})^2 - \sqrt{t} - 40 = 0$$

$$\sqrt{t} = \frac{1 \pm \sqrt{1 + 320}}{4} = 4.73$$

$$t = 22.4$$

round up to 23

Test Question: 107

Key: C

X = losses on one life

$$\begin{aligned} E[X] &= (0.3)(1) + (0.2)(2) + (0.1)(3) \\ &= 1 \end{aligned}$$

S = total losses

$$E[S] = 3E[X] = 3$$

$$\begin{aligned} E[(S-1)_+] &= E[S] - 1(1 - F_s(0)) \\ &= E[S] - (1)(1 - f_s(0)) \\ &= 3 - (1)(1 - 0.4^3) \\ &= 3 - 0.936 \\ &= 2.064 \end{aligned}$$

Test Question: 108

Key: C

$$\begin{aligned} p(k) &= \frac{2}{k} p(k-1) \\ &= \left[0 + \frac{2}{k} \right] p(k-1) \end{aligned}$$

Thus an $(a, b, 0)$ distribution with $a = 0, b = 2$.

Thus Poisson with $\lambda = 2$.

$$\begin{aligned} p(4) &= \frac{e^{-2} 2^4}{4!} \\ &= 0.09 \end{aligned}$$

Test Question: 109

Key: B

By the memoryless property, the distribution of amounts paid in excess of 100 is still exponential with mean 200.

With the deductible, the probability that the amount paid is 0 is

$$F(100) = 1 - e^{-100/200} = 0.393.$$

Thus the average amount paid per loss is $(0.393)(0) + (0.607)(200) = 121.4$

The expected number of losses is $(20)(0.8) = 16$.

The expected amount paid is $(16)(121.4) = 1942$.

Test Question: 110**Key: E**

$$E[N] = (0.8)(1) + (0.2)(2) = 1.2$$

$$E[N^2] = (0.8)1 + (0.2)(4) = 1.6$$

$$\text{Var}(N) = 1.6 - 1.2^2 = 0.16$$

$$E[X] = 70 + 100 = 170$$

$$\text{Var}(X) = E[X^2] - E[X]^2 = (7000 + 100,000) - 170^2 = 78,100$$

$$E[S] = E[N]E[X] = 1.2(170) = 204$$

$$\text{Var}(S) = E[N]\text{Var}(X) + E[X]^2\text{Var}(N) = 1.2(78,100) + 170^2(0.16) = 98,344$$

$$\text{Std dev } (S) = \sqrt{98,344} = 313.6$$

$$\text{So } B = 204 + 314 = 518$$

Test Question: 111**Key: E**

Treat as three independent Poisson variables, corresponding to 1, 2 or 3 claimants.

$$\text{rate}_1 = 6 \left[= \frac{1}{2} \times 12 \right]$$

$$\text{rate}_2 = 4$$

$$\text{rate}_3 = 2$$

$$\text{Var}_1 = 6$$

$$\text{Var}_2 = 16 \left[= 4 \times 2^2 \right]$$

$$\text{Var}_3 = 18$$

total Var = 6 + 16 + 18 = 40, since independent.

Alternatively,

$$E(X^2) = \frac{1^2}{2} + \frac{2^2}{3} + \frac{3^2}{6} = \frac{10}{3}$$

$$\begin{aligned} \text{For compound Poisson, } \text{Var}[S] &= E[N]E[X^2] \\ &= (12)\left(\frac{10}{3}\right) = 40 \end{aligned}$$

Test Question: 112**Key: A**

N = number of physicians $E(N) = 3$ $\text{Var}(N) = 2$
 X = visits per physician $E(X) = 30$ $\text{Var}(X) = 30$
 S = total visits

$$E(S) = E(N) E(X) = 90$$

$$\text{Var}(S) = E(N) \text{Var}(X) + E^2(X) \text{Var}(N) =$$

$$= 3 \cdot 30 + 900 \cdot 2 = 1890$$

Standard deviation (S) = 43.5

$$\Pr(S > 119.5) = \Pr\left(\frac{S - 90}{43.5} > \frac{119.5 - 90}{43.5}\right) = 1 - \Phi(0.68) \text{ Course 3: November 2000}$$

Test Question: 113**Key: E**

$$E(N) = .7$$

$$\text{Var}(N) = 4 \times .2 + 9 \times .1 - .49 = 1.21$$

$$E(X) = 2$$

$$\text{Var}(X) = 100 \times .2 - 4 = 16$$

$$E(S) = 2 \times .7 = 1.4$$

$$\text{Var}(S) = E(N) \text{Var}(X) + E^2(X) \text{Var}(N) =$$

$$= .7 \times 16 + 4 \times 1.21 = 16.04$$

Standard Dev(S) = 4

$$E(S) + 2 \times \text{Standard Dev}(S) = 1.4 + 2 \times 4 = 9.4$$

Since there are no possible values of S between 0 and 10,

$$\Pr(S > 9.4) = 1 - \Pr(S = 0)$$

$$= 1 - .7 - .2 \times .8^2 - .1 \times .8^3 = .12$$

Test Question: 114**Key: A**

$$P(0) = \frac{1}{5} \int_0^5 e^{-\lambda} d\lambda = \frac{1}{5} (-e^{-\lambda}) \Big|_0^5 = \frac{1}{5} (1 - e^{-5}) = 0.1987$$

$$P(1) = \frac{1}{5} \int_0^5 \lambda e^{-\lambda} d\lambda = \frac{1}{5} (-\lambda e^{-\lambda} - e^{-\lambda}) \Big|_0^5 = \frac{1}{5} (1 - 6e^{-5}) = 0.1919$$

$$P(N \geq 2) = 1 - .1987 - .1919 = .6094$$

Test Question: 115**Key: D**

Let X be the occurrence amount, $Y = \max(X-100, 0)$ be the amount paid.

$$E[X] = 1,000$$

$$\text{Var}[X] = (1,000)^2$$

$$P(X > 100) = \exp(-100/1,000) = .904837$$

The distribution of Y given that $X > 100$, is also exponential with mean 1,000 (memoryless property).

So Y is $\begin{cases} 0 \text{ with prob } .095163 \\ \text{exponential mean } 1000 \text{ with prob } .904837 \end{cases}$

$$E[Y] = .095163 \times 0 + .904837 \times 1,000 = 904.837$$

$$E[Y^2] = .095163 \times 0 + .904837 \times 2 \times (1,000)^2 = 1,809,675$$

$$\text{Var}[Y] = 1,809,675 - (904.837)^2 = 990,944$$

Alternatively, think of this as a compound distribution whose frequency is Bernoulli with $p = .904837$, and severity is exponential with mean 1,000.

$$\text{Var} = \text{Var}[N] \times E[X]^2 + \text{Var}[X] \times E[N] = p(1-p)(1,000,000) + p(1,000,000)$$

Test Question: 116**Key: C**

Expected claims under current distribution = 500

θ = parameter of new distribution

X = claims

$$E(X) = \theta$$

$$\text{bonus} = .5 \times [500 - X \wedge 500]$$

$$E(\text{claims} + \text{bonus}) = \theta + .5 \left(500 - \theta \left(1 - \frac{\theta}{500 + \theta} \right) \right) = 500$$

$$\theta - \frac{\theta}{2} \left(\frac{500}{500 + \theta} \right) = 250$$

$$2(500 + \theta)\theta - 500\theta = 250(500 + \theta) \cdot 2$$


$$1000\theta + \theta^2 \cdot 2 - 500\theta = 2 \times 250 \times 500 + 500\theta$$

$$\theta = \sqrt{250 \times 500} = 354$$

Test Question: 117**Key: A**

$$UDD \Rightarrow l_{21} = (0.8)(53,488) + (0.2)(17,384) = 46,267.2$$

$$\begin{aligned} Mrl(21) &= \overset{\circ}{e}_{21} = \int_0^{\infty} {}_t p_{21} \int_0^{\infty} \frac{S(21+t)}{S(21)} dt \\ &= \sum \text{areas} \\ &= 2.751 + 1.228 + 0.361 + 0.072 \\ &= 4.412 \end{aligned}$$

**Test Question: 118****Key: D**

$$\mu'_{n1} = E[N] = 25$$

$$\mu_{x2} = \text{Var}[X] = 675$$

$$\mu_{N2} = \text{Var}[N] = 25$$

$$E[X] = 50$$

$$E[S] = E[X]E[N] = 25 \times 50 = 1250$$

$$\begin{aligned} \text{Var}[S] &= E[N]\text{Var}[X] + \text{Var}[N]E[X]^2 \\ &= 25 \times 675 + 25 \times 2500 = 79,375 \end{aligned}$$

$$\text{Standard Deviation}[S] = \sqrt{79,375} = 281.74$$

$$\Pr(S > 2000) = \Pr\left[\frac{(S - 1250)}{281.74} > \frac{(2000 - 1250)}{281.74}\right] = 1 - \Phi(2.66)$$

Test Question: 119

Key: C

$$E[X] = 2000(1!) / (1!) = 2000$$

$$E[X \wedge 3000] = \left(\frac{2000}{1}\right) \times \left[1 - \frac{2000}{(3000 + 2000)}\right] = 2000 \times \left(1 - \frac{2}{5}\right) = 2000 \times \frac{3}{5} = 1200$$

So the fraction of the losses expected to be covered by the reinsurance is $\frac{2000 - 1200}{2000} = 0.4$.

The expected ceded losses are 4,000,000 \Rightarrow the ceded premium is 4,400,000.

Test Question: 120

Key: E

$$X_{2002} = 1.05 \times X_{2001}$$

$$\begin{aligned} \text{so: } F\left(\frac{x_{2002}}{1.05}\right) &= 1 - \left[\frac{2000}{(x_{2002} / 1.05 + 2000)}\right]^2 \\ &= 1 - \left[\frac{2100}{x_{2002} + 2100}\right]^2 \end{aligned}$$

This is just another Pareto distribution with $\alpha = 2, \theta = 2100$.

$$E[X_{2002}] = 2100.$$

and

$$E[X_{2002} \wedge 3000] = \left(\frac{2100}{1}\right) \times \left[1 - \left(\frac{2100}{(3000 + 2100)}\right)\right]$$

$$= 2100 \times \left[\frac{3000}{5100}\right] = 1235$$

So the fraction of the losses expected to be covered by the reinsurance is

$$\frac{2100 - 1235}{2100} = 0.412.$$

The total expected losses have increased to 10,500,000, so

$$C_{2002} = 1.1 \times 0.412 \times 10,500,000 = 4,758,600$$

And $\frac{C_{2002}}{C_{2001}} = \frac{4,758,600}{4,400,000} = 1.08$

Question #121
DELETED

Question #122
Key: C

$$E[N] = mq = 1.8 \Rightarrow q = \frac{1.8}{3} = 0.6$$

x	$f_N(x)$	$F_N(x)$
0	0.064	0.064
1	0.288	0.352
2	0.432	0.784
3	0.216	1.000

First: $0.432 < 0.7 < 0.784$ so $N = 2$.

Second: $0.064 < 0.1 < 0.352$ so $N = 1$

Third: $0.432 < 0.5 < 0.784$ so $N = 2$

Use 0.1 and 0.3 for amounts

Use 0.9 for amount

Use 0.5 and 0.7 for amounts

Discrete uniform $\Rightarrow F_X(x) = 0.2x, \quad x = 1, 2, 3, 4, 5$

$$0.4 < 0.5 < 0.6 \Rightarrow x_1 = 3$$

$$0.6 < 0.7 < 0.8 \Rightarrow x_2 = 4$$

Aggregate claims = $3+4 = 7$

Question #123**Key: C**

$$E(X \wedge x) = \frac{\theta}{\alpha - 1} \left[1 - \left(\frac{\theta}{x + \theta} \right)^{\alpha - 1} \right] = \frac{2000x}{x + 2000}$$

X	$E(X \wedge x)$
∞	2000
250	222
2250	1059
5100	1437

$$0.75(E(X \wedge 2250) - E(X \wedge 250)) + 0.95(E(X) - E(X \wedge 5100))$$

$$0.75(1059 - 222) + 0.95(2000 - 1437) = 1162.6$$

The 5100 breakpoint was determined by when the insured's share reaches 3600:

$$3600 = 250 + 0.25(2250 - 250) + (5100 - 2250)$$

Question #124**Key: D**

Since each time the probability of a heavy scientist is just half the probability of a success, the distribution is binomial with $q = 0.6 \times 0.5 = 0.3$ and $m = 8$.

$$f(2) = \binom{8}{2} \times (0.3)^2 \times (0.7)^6 = 0.30$$

Question #125**Key: A**

Let N_1, N_2 denote the random variable for # of claims for Type I and II in 2 years

X_1, X_2 denote the claim amount for Type I and II

$S_1 =$ total claim amount for type I in 2 years

$S_2 =$ total claim amount for Type II at time in 2 years

$S = S_1 + S_2 =$ total claim amount in 2 years

$\{S_1\} \rightarrow$ compound poisson $\lambda_1 = 2 \times 6 = 12$ $X_1 \sim U(0, 1)$

$\{S_2\} \rightarrow$ compound poisson $\lambda_2 = 2 \times 2 = 4$ $X_2 \sim U(0, 5)$

$$E(N_1) = \text{Var}(N_1) = 2 \times 6 = 12$$

$$E(S_1) = E(N_1)E(X_1) = (12)(0.5) = 6$$

$$\begin{aligned}\text{Var}(S_1) &= E(N_1)\text{Var}(X_1) + \text{Var}(N_1)(E(X_1))^2 \\ &= (12)\frac{(1-0)}{12} + (12)(0.5)^2 \\ &= 4\end{aligned}$$

$$E(N_2) = \text{Var}(N_2) = 2 \times 2 = 4$$

With formulas corresponding to those for S_1 ,

$$E(S_2) = 4 \times \frac{5}{2} = 10$$

$$\text{Var}(S_2) = 4 \times \frac{(5-0)^2}{12} + 4\left(\frac{5}{2}\right)^2 = 33.\bar{3}$$

$$E(S) = E(S_1) + E(S_2) = 6 + 10 = 16$$

Since S_1 and S_2 are independent,

$$\text{Var}(S) = \text{Var}(S_1) + \text{Var}(S_2) = 4 + 33.\bar{3} = 37.\bar{3}$$

$$\Pr(S > 18) = \Pr\left(\frac{S-16}{\sqrt{39.\bar{3}}} > \frac{2}{\sqrt{37.\bar{3}}} = 0.327\right)$$

Using normal approximation

$$\begin{aligned}\Pr(S > 18) &= 1 - \Phi(0.327) \\ &= 0.37\end{aligned}$$

Question #126

Key: C

Let X be the loss random variable,

So $(X - 5)_+$ is the claim random variable.

$$E(X) = \frac{10}{2.5-1} = 6.\bar{6}$$

$$E(X \wedge 5) = \left(\frac{10}{2.5-1} \right) \left[1 - \left(\frac{10}{5+10} \right)^{2.5-1} \right]$$

$$= 3.038$$

$$E(X-5)_+ = E(X) - E(X \wedge 5)$$

$$= 6.6 - 3.038$$

$$= 3.629$$

$$\text{Expected aggregate claims} = E(N)E(X-5)_+$$

$$= (5)(3.629)$$

$$= 18.15$$

Question #127

Key: B

A Pareto ($\alpha = 2, \theta = 5$) distribution with 20% inflation becomes Pareto with

$$\alpha = 2, \theta = 5 \times 1.2 = 6$$

$$\text{In 2004, } E(X) = \frac{6}{2-1} = 6$$

$$E(X \wedge 10) = \frac{6}{2-1} \left(1 - \left(\frac{6}{10+6} \right)^{2-1} \right) = 3.75$$

$$E(X-10)_+ = E(X) - E(X \wedge 10)$$

$$= 6 - 3.75 = 2.25$$

$$\text{LER} = 1 - \frac{E(X-10)_+}{E(X)} = 1 - \frac{2.25}{6} = 0.625$$

Question #128

DELETED

Question #129
DELETED

Question #130
Key: E

$$\begin{aligned} E(W) &= \frac{1}{4} \int_0^4 \sum_{i=0}^{\infty} 2^i \Pr(N = i | \lambda) d\lambda \quad \left[\frac{1}{4} \text{ is the density of } \lambda \text{ on } [0, 4]. \right] \\ &= \frac{1}{4} \int_0^4 P(2 | \lambda) d\lambda \quad [\text{see note}] \\ &= \frac{1}{4} \int_0^4 e^{\lambda(2-1)} d\lambda \quad [\text{using formula from tables for the pgf of the Poisson}] \\ &= \frac{1}{4} e^{\lambda} \Big|_0^4 = \frac{1}{4} (e^4 - 1) \\ &= 13.4 \end{aligned}$$

Note: the probability generating function (pgf) is $P(Z) = \sum_{k=0}^{\infty} p_k Z^k$ so the integrand is $P(2)$, or in this case $P(2 | \lambda)$ since λ is not known.

Alternatively,

$$\begin{aligned} E(W) &= \frac{1}{4} \int_0^4 \sum_{i=0}^{\infty} 2^i \Pr(N = i | \lambda) d\lambda \\ &= \frac{1}{4} \int_0^4 \sum_{i=0}^{\infty} \frac{2^i e^{-\lambda} \lambda^i}{i!} d\lambda \\ &= \frac{1}{4} \int_0^4 \sum_{i=0}^{\infty} \frac{e^{-\lambda} (2\lambda)^i}{i!} d\lambda \end{aligned}$$

We know $\sum_{i=0}^{\infty} \frac{e^{-2\lambda} (2\lambda)^i}{i!} = 1$ since $\frac{e^{-2\lambda} (2\lambda)^i}{i!}$ is $f(i)$ for a Poisson with mean 2λ

$$\text{so } \sum_{i=0}^{\infty} \frac{e^{-\lambda} (2\lambda)^i}{i!} = \frac{e^{-\lambda}}{e^{-2\lambda}} = e^{\lambda}$$

Thus $E(W) = \frac{1}{4} \int_0^4 e^{\lambda} d\lambda$

$$= \frac{1}{4} e^{\lambda} \Big|_0^4 = \frac{1}{4} (e^4 - 1)$$

$$= 13.4$$

Question #131

Key: E

$$E(S) = \lambda E[X] = 2/3(1/4 + 2/4 + 3/2) = 2/3 \times 9/4 = 3/2$$

$$\text{Var}(S) = \lambda E[X^2] = 2/3(1/4 + 4/4 + 9/2) = 23/6$$

So cumulative premium to time 2 is $2(3/2 + 1.8\sqrt{23/6}) = 10$, where the expression in parentheses is the annual premium

Times between claims are determined by $-(1/\lambda) \log(1-u)$ and are 0.43, 0.77, 1.37, 2.41

So 2 claims before time 2 (second claim is at 1.20; third is at 2.57)

Sizes are 2, 3, 1, 3, where only the first two matter.

So gain to the insurer is $10 - (2+3) = 5$

Question #132

Key: C

To get number of claims, set up cdf for Poisson:

x	$f(x)$	$F(x)$
0	0.135	0.135
1	0.271	0.406
2	0.271	0.677
3	0.180	0.857

0.80 simulates 3 claims.

$$F(x) = 1 - (500/(x+500))^2 = u, \text{ so } x = (1-u)^{-1/2} 500 - 500$$

0.6 simulates 290.57

0.25 simulates 77.35
0.7 simulates 412.87

So total losses equals 780.79

Insurer pays $(0.80)(750) + (780.79 - 750) = 631$

Question #133

Key: C

$$E(X | q) = 3q, \text{Var}(X | q) = 3q(1 - q)$$

$$\mu = E(3q) = \int_0^1 3q \cdot 2q dq = 2q^3 \Big|_0^1 = 2$$

$$v = E[3q(1 - q)] = \int_0^1 3q(1 - q) \cdot 2q dq = 2q^3 - 1.5q^4 \Big|_0^1 = 0.5$$

$$a = \text{Var}(3q) = E(9q^2) - \mu^2 = \int_0^1 9q^2 \cdot 2q dq - 2^2 = 4.5q^4 \Big|_0^1 - 4 = 4.5 - 4 = 0.5$$

$$k = v/a = 0.5/0.5 = 1$$

$$Z = \frac{1}{1+1} = 0.5$$

The estimate is

$$0.5(0) + 0.5(2) = 1.$$

Question #134

Key: D

$$0.35(14) = 4.9$$

$$\hat{\pi}_{0.35} = 0.1(216) + 0.9(250) = 246.6$$

Question #135

Key: E

At $y_1 = 0.9$ the risk set is $r_1 = 7$ and $s_1 = 1$.

At $y_2 = 1.5$ the risk set is $r_2 = 6$ and $s_2 = 1$.

$$\text{Then, } S_{10}(1.6) = \frac{6}{7} \frac{5}{6} = 0.7143.$$

Question #136

Key: B

$$\Pr(\text{class1} | \text{claim} = 250) = \frac{\Pr(\text{claim} = 250 | \text{class1}) \Pr(\text{class1})}{\Pr(\text{claim} = 250 | \text{class1}) \Pr(\text{class1}) + \Pr(\text{claim} = 250 | \text{class2}) \Pr(\text{class2})}$$

$$= \frac{0.5(2/3)}{0.5(2/3) + 0.7(1/3)} = \frac{10}{17}.$$

$$E(\text{claim} | \text{class1}) = 0.5(250) + 0.3(2,500) + 0.2(60,000) = 12,875.$$

$$E(\text{claim} | \text{class2}) = 0.7(250) + 0.2(2,500) + 0.1(60,000) = 6,675.$$

$$E(\text{claim} | 250) = \frac{10}{17}(12,875) + \frac{7}{17}(6,675) = 10,322.$$

Question #137

Key: D

$$L(p) = f(0.74)f(0.81)f(0.95)$$

$$= (p+1)0.74^p (p+1)0.81^p (p+1)0.95^p$$

$$= (p+1)^3(0.56943)^p$$

$$l(p) = \ln L(p) = 3\ln(p+1) + p\ln(0.56943)$$

$$l'(p) = \frac{3}{p+1} - 0.563119 = 0$$

$$p+1 = \frac{3}{0.563119} = 5.32747$$

$$p = 4.32747.$$

Question #138

Key: E

The sample mean is 1 and therefore $mq = 1$.

For the smoothed empirical 33rd percentile, $(1/3)(5 + 1) = 2$ and the second smallest sample item is 0. For the 33rd percentile of the binomial distribution to be 0, the probability at zero must exceed 0.33. So, $(1-q)^m > 0.33$ and then $(1-m^{-1})^m > 0.33$. Trial and error gives $m = 6$ as the smallest value that produces this result.

Question #139

Key: C

Let X be the number of claims.

$$E(X | I) = 0.9(0) + 0.1(2) = 0.2$$

$$E(X | II) = 0.8(0) + 0.1(1) + 0.1(2) = 0.3$$

$$E(X | III) = 0.7(0) + 0.2(1) + 0.1(2) = 0.4$$

$$\text{Var}(X | I) = 0.9(0) + 0.1(4) - 0.2^2 = 0.36$$

$$\text{Var}(X | II) = 0.8(0) + 0.1(1) + 0.1(4) - 0.3^2 = 0.41$$

$$\text{Var}(X | III) = 0.7(0) + 0.2(1) + 0.1(4) - 0.4^2 = 0.44.$$

$$\mu = (1/2)(0.2 + 0.3 + 0.4) = 0.3$$

$$v = (1/3)(0.36 + 0.41 + 0.44) = 0.403333$$

$$a = (1/3)(0.2^2 + 0.3^2 + 0.4^2) - 0.3^2 = 0.006667$$

$$k = 0.403333 / 0.006667 = 60.5$$

$$Z = \frac{50}{50 + 60.5} = 0.45249.$$

For one insured the estimate is $0.45249(17/50) + 0.54751(0.3) = 0.31810$.

For 35 insureds the estimate is $35(0.31810) = 11.13$.

Question #140

Key: A

For the given intervals, based on the model probabilities, the expected counts are 4.8, 3.3, 8.4, 7.8, 2.7, 1.5, and 1.5. To get the totals above 5, group the first two intervals and the last three. The table is

Interval	Observed	Expected	Chi-square
0—500	3	8.1	3.21
500—2498	8	8.4	0.02
2498—4876	9	7.8	0.18
4876—infinity	10	5.7	3.24
Total	30	30	6.65

Question #141

Key: E

Let $\hat{H} = \hat{H}(t)$ and $\hat{v} = \text{Var}(\hat{H}(t))$. The confidence interval is $\hat{H}U$ where

$U = \exp(\pm z_{\alpha/2} \sqrt{\hat{v}} / \hat{H})$. Multiplying the two bounds gives

$0.7(0.357) = \hat{H}^2$ for $\hat{H} = 0.49990$. Then,

$$\hat{S} = \exp(-0.49990) = 0.60659.$$

Question #142

Key: C

$$\begin{aligned}
0.575 &= \Pr(N = 0) = \int_0^k \Pr(N = 0 | \theta) \pi(\theta) d\theta \\
&= \int_0^k e^{-\theta} \frac{e^{-\theta}}{1 - e^{-k}} d\theta = -\frac{e^{-2\theta}}{2(1 - e^{-k})} \Big|_0^k = -\frac{e^{-2k}}{2(1 - e^{-k})} + \frac{1}{2(1 - e^{-k})} \\
&= \frac{1 - e^{-2k}}{2(1 - e^{-k})} = \frac{1 + e^{-k}}{2} \\
e^{-k} &= 2(0.575) - 1 = 0.15 \\
k &= 1.90.
\end{aligned}$$

Question #143**Key: C**

The sample -1 moment is $\frac{1}{6}\left(\frac{1}{15} + \frac{1}{45} + \frac{1}{140} + \frac{1}{250} + \frac{1}{560} + \frac{1}{1340}\right) = 0.017094$. The sample -

2 moment is $\frac{1}{6}\left(\frac{1}{15^2} + \frac{1}{45^2} + \frac{1}{140^2} + \frac{1}{250^2} + \frac{1}{560^2} + \frac{1}{1340^2}\right) = 0.00083484$.

Then the equations are

$$0.017094 = \frac{1}{\theta(\tau-1)},$$

$$0.00083484 = \frac{2}{\theta^2(\tau-1)(\tau-2)}.$$

Divide the square of the first equation by the second equation to obtain

$$0.35001 = \frac{\tau-2}{2(\tau-1)} \text{ which is solved for } \tau = 4.33356. \text{ From the first equation,}$$

$$\theta = \frac{1}{3.33356(0.017094)} = 17.55.$$

Question #144**Key: A**

For each simulation, estimate the LER and then calculate the squared difference from the estimate, 0.125.

Simulation	First claim	Second claim	Third claim	LER	Squared difference
1	600	600	1500	0.111111	0.000193
2	1500	300	1500	0.090909	0.001162
3	1500	300	600	0.125000	0.000000
4	600	600	300	0.200000	0.005625
5	600	300	1500	0.125000	0.000000
6	600	600	1500	0.111111	0.000193
7	1500	1500	1500	0.066667	0.003403
8	1500	300	1500	0.090909	0.001162
9	300	600	300	0.250000	0.015625
10	600	600	600	0.166667	0.001736

The last column has an average of 0.002910 which is the bootstrap estimate.

Question #145**Key: B**

The subscripts denote the three companies.

$$\begin{aligned}
x_{I1} &= \frac{50,000}{100} = 500, & x_{I2} &= \frac{50,000}{200} = 250, & x_{III} &= \frac{150,000}{500} = 300 \\
x_{II2} &= \frac{150,000}{300} = 500, & x_{III1} &= \frac{150,000}{50} = 3,000, & x_{III2} &= \frac{150,000}{150} = 1,000 \\
\bar{x}_I &= \frac{100,000}{300} = 333.33, & \bar{x}_{II} &= \frac{300,000}{800} = 375, & \bar{x}_{III} &= \frac{300,000}{200} = 1,500, & \bar{x} &= \frac{700,000}{1,300} = 538.46 \\
\hat{v} &= \frac{100(500 - 333.33)^2 + 200(250 - 333.33)^2 + 500(300 - 375)^2 + 300(500 - 375)^2}{(2-1) + (2-1) + (2-1)} \\
&= 53,888,888.89 \\
\hat{a} &= \frac{300(333.33 - 538.46)^2 + 800(375 - 538.46)^2 + 200(1,500 - 538.46)^2 - 53,888,888.89(3-1)}{1,300 - \frac{300^2 + 800^2 + 200^2}{1,300}} \\
&= 157,035.60 \\
k &= \frac{53,888,888.89}{157,035.60} = 343.1635, & Z &= \frac{200}{200 + 343.1635} = 0.3682
\end{aligned}$$

Question #146

Key: D

Let α_j be the parameter for region j . The likelihood function is $L = \left(\prod_{i=1}^n \frac{\alpha_1}{x_i^{\alpha_1+1}} \right) \left(\prod_{i=1}^m \frac{\alpha_2}{y_i^{\alpha_2+1}} \right)$.

The expected values satisfy $\frac{\alpha_2}{\alpha_2 - 1} = 1.5 \frac{\alpha_1}{\alpha_1 - 1}$ and so $\alpha_2 = \frac{3\alpha_1}{2 + \alpha_1}$. Substituting this in the likelihood function and taking logs produces

$$l(\alpha_1) = \ln L(\alpha_1) = n \ln \alpha_1 - (\alpha_1 + 1) \sum_{i=1}^n \ln x_i + m \ln \left(\frac{3\alpha_1}{2 + \alpha_1} \right) - \frac{2 + 4\alpha_1}{2 + \alpha_1} \sum_{i=1}^m \ln y_i$$

$$l'(\alpha_1) = \frac{n}{\alpha_1} - \sum_{i=1}^n \frac{1}{x_i} + \frac{2m}{\alpha_1(2 + \alpha_1)} - \frac{6}{(2 + \alpha_1)^2} \sum_{i=1}^m \ln y_i = 0.$$

Question #147**Key: D**

Let $K_y(x)$ be the contribution at x of the data point at y . It is

$$K_y(x) = \begin{cases} 0, & x < y - 1.4 \\ \frac{x - y + 1.4}{2.8}, & y - 1.4 \leq x \leq y + 1.4 \\ 1, & x > y + 1.4. \end{cases}$$

For the particular points,

$K_2(4) = 1$, $K_{3.3}(4) = \frac{4 - 3.3 + 1.4}{2.8} = 0.75$, $K_4(4) = 0.5$, $K_{4.7}(4) = 0.25$. The kernel estimate is the weighted average $\frac{1}{8}(1) + \frac{2}{8}(0.75) + \frac{2}{8}(0.5) + \frac{3}{8}(0.25) = 0.53125$.

Question #148**Key: B**

The mean is mq and the variance is $mq(1 - q)$. The mean is 34,574 and so the full credibility standard requires the confidence interval to be ± 345.74 which must be 1.96 standard deviations. Thus,

$$345.74 = 1.96\sqrt{mq(1 - q)} = 1.96\sqrt{34,574}\sqrt{1 - q}$$

$$1 - q = 0.9, \quad q = 0.1.$$

Question #149**Key: A**

Only the Kolmogorov-Smirnov test statistic tends toward zero as the sample size goes to infinity. As a consequence, the critical value for the K-S statistic has the square root of the sample size in the denominator. For the Anderson-Darling and the Chi-square goodness-of-fit test statistics, the sample size appears in the numerator of the test statistics themselves. The Schwarz Bayesian Criterion involves an adjustment to the likelihood function, which does not go to zero as the sample size goes to infinity.

Question #150**Key: E**

The sample average is $(14 + 33 + 72 + 94 + 120 + 135 + 150 + 150)/8 = 96$. The model

average is $E(X \wedge 150) = \int_0^{150} x \frac{1}{\theta} dx + \int_{150}^{\theta} 150 \frac{1}{\theta} dx = \frac{150^2}{2\theta} + 150 \frac{\theta - 150}{\theta} = 150 - \frac{11,250}{\theta}$. The

equation to solve is $150 - \frac{11,250}{\theta} = 96$, $\frac{11,250}{\theta} = 54$, $\theta = \frac{11,250}{54} = 208.3$.

Question #151**Key: C**

$$E(N | 1) = 5, E(N | 2) = 8(0.55) = 4.4, \mu = 0.5(5) + 0.5(4.4) = 4.7$$

$$\text{Var}(N | 1) = 5, \text{Var}(N | 2) = 8(0.55)(0.45) = 1.98, v = 0.5(5) + 0.5(1.98) = 3.49$$

$$a = 0.5(5)^2 + 0.5(4.4)^2 - 4.7^2 = 0.09, k = 3.49/0.09 = 38.7778$$

$$Z = \frac{3}{3 + 38.7778} = 0.0718, 4.6019 = 0.0718 \frac{7+r}{3} + 0.9282(4.7)$$

The solution is $r = 3$.

Question #152**Key: A**

These observations are truncated at 500. The contribution to the likelihood function is

$$\frac{f(x)}{1 - F(500)} = \frac{\theta^{-1} e^{-x/\theta}}{e^{-500/\theta}}. \text{ Then the likelihood function is}$$

$$L(\theta) = \frac{\theta^{-1} e^{-600/\theta} \theta^{-1} e^{-700/\theta} \theta^{-1} e^{-900/\theta}}{(e^{-500/\theta})^3} = \theta^{-3} e^{-700/\theta}$$

$$l(\theta) = \ln L(\theta) = -3 \ln \theta - 700\theta^{-1}$$

$$l'(\theta) = -3\theta^{-1} + 700\theta^{-2} = 0$$

$$\theta = 700/3 = 233.33.$$

Question #153
DELETED

Question #154
Key: E

For a compound Poisson distribution, S , the mean is $E(S | \lambda, \mu, \sigma) = \lambda E(X) = \lambda e^{\mu+0.5\sigma^2}$ and the variance is $Var(S | \lambda, \mu, \sigma) = \lambda E(X^2) = \lambda e^{2\mu+2\sigma^2}$. Then,

$$\begin{aligned} E(S) &= E[E(S | \lambda, \mu, \sigma)] = \int_0^1 \int_0^1 \int_0^1 \lambda e^{\mu+0.5\sigma^2} 2\sigma d\lambda d\mu d\sigma \\ &= \int_0^1 \int_0^1 e^{\mu+0.5\sigma^2} \sigma d\mu d\sigma = \int_0^1 (e-1)e^{0.5\sigma^2} \sigma d\sigma \\ &= (e-1)(e^{0.5} - 1) = 1.114686 \end{aligned}$$

$$\begin{aligned} v &= E[Var(S | \lambda, \mu, \sigma)] = \int_0^1 \int_0^1 \int_0^1 \lambda e^{2\mu+2\sigma^2} 2\sigma d\lambda d\mu d\sigma \\ &= \int_0^1 \int_0^1 e^{2\mu+2\sigma^2} \sigma d\mu d\sigma = \int_0^1 0.5(e^2 - 1)e^{2\sigma^2} \sigma d\sigma \\ &= 0.5(e^2 - 1)0.25(e^2 - 1) = 0.125(e^2 - 1)^2 = 5.1025 \end{aligned}$$

$$\begin{aligned} a &= Var[E(S | \lambda, \mu, \sigma)] = \int_0^1 \int_0^1 \int_0^1 \lambda^2 e^{2\mu+\sigma^2} 2\sigma d\lambda d\mu d\sigma - E(S)^2 \\ &= \int_0^1 \int_0^1 \frac{2}{3} e^{2\mu+\sigma^2} \sigma d\mu d\sigma - E(S)^2 = \int_0^1 \frac{1}{3} (e^2 - 1)e^{\sigma^2} \sigma d\sigma - E(S)^2 \\ &= \frac{1}{3}(e^2 - 1)\frac{1}{2}(e - 1) - E(S)^2 = (e^2 - 1)(e - 1)/6 - E(S)^2 = 0.587175 \end{aligned}$$

$$k = \frac{5.1025}{0.587175} = 8.69.$$

Question #155
Key: D

The equations to solve are $0.4 = e^{-(\theta/1.82)^\tau}$, $0.8 = e^{-(\theta/12.66)^\tau}$. Taking logs yields $0.91629 = (\theta/1.82)^\tau$, $0.22314 = (\theta/12.66)^\tau$. Taking the ratio of the first equation to the second equation gives $4.10635 = (12.66/1.82)^\tau = 6.95604^\tau$. Taking logs again, $1.41253 = 1.93961\tau$ and then $\tau = 0.72825$. Returning to the first (logged) equation, $0.91629 = (\theta/1.82)^\tau$, $0.88688 = \theta/1.82$, $\theta = 1.614$.

Question #156**Key: C**

There are $n/2$ observations of $N = 0$ (given $N = 0$ or 1) and $n/2$ observations of $N = 1$ (given $N = 0$ or 1). The likelihood function is

$$L = \left(\frac{e^{-\lambda}}{e^{-\lambda} + \lambda e^{-\lambda}} \right)^{n/2} \left(\frac{\lambda e^{-\lambda}}{e^{-\lambda} + \lambda e^{-\lambda}} \right)^{n/2} = \frac{\lambda^{n/2} e^{-n\lambda}}{(e^{-\lambda} + \lambda e^{-\lambda})^n} = \frac{\lambda^{n/2}}{(1 + \lambda)^n}.$$

Taking logs, differentiating and solving provides the answer.

$$l = \ln L = (n/2) \ln \lambda - n \ln(1 + \lambda)$$

$$l' = \frac{n}{2\lambda} - \frac{n}{1 + \lambda} = 0$$

$$n(1 + \lambda) - n2\lambda = 0$$

$$1 - \lambda = 0, \quad \lambda = 1.$$

Question #157**Key: D**

The posterior density function is proportional to the product of the likelihood function and prior density. That is, $\pi(q|1,0) \propto f(1|q)f(0|q)\pi(q) \propto q(1-q)q^3 = q^4 - q^5$. To get the exact posterior density, integrate this function over its range:

$$\int_{0.6}^{0.8} q^4 - q^5 dq = \left. \frac{q^5}{5} - \frac{q^6}{6} \right|_{0.6}^{0.8} = 0.014069 \text{ and so } \pi(q|1,0) = \frac{q^4 - q^5}{0.014069}.$$

$$\Pr(0.7 < q < 0.8 | 1,0) = \int_{0.7}^{0.8} \frac{q^4 - q^5}{0.014069} dq = 0.5572.$$

Question #158**Key: B**

The cumulative hazard function for the exponential distribution is $H(x) = x/\theta$. The maximum likelihood estimate of θ is the sample mean, which equals $(1227/15) = 81.8$. Therefore $\hat{H}_2(75) = (75/81.8) = 0.917$.

To calculate $\hat{H}_1(75)$ use the following table.

j	1	2	3	4	5	6
y_j	11	22	36	51	69	92
s_j	1	3	1	1	3	2
r_j	15	14	11	10	9	6

Therefore,

$$\hat{H}_1(75) = \frac{1}{15} + \frac{3}{14} + \frac{1}{11} + \frac{1}{10} + \frac{3}{9} = 0.805.$$

Thus, $\hat{H}_2(75) - \hat{H}_1(75) = 0.917 - 0.805 = 0.112$.

Question #159

Key: A

The sample mean is $\frac{0(2000) + 1(600) + 2(300) + 3(80) + 4(20)}{3000} = 0.5066667 = \hat{\mu} = \hat{v}$ and the

sample variance is

$$\frac{2000(0 - \hat{\mu})^2 + 600(1 - \hat{\mu})^2 + 300(2 - \hat{\mu})^2 + 80(3 - \hat{\mu})^2 + 20(4 - \hat{\mu})^2}{2999} = 0.6901856. \text{ Then,}$$

$$\hat{a} = 0.6901856 - 0.5066667 = 0.1835189, k = \frac{0.5066667}{0.1835189} = 2.760842 \text{ and}$$

$$Z = \frac{1}{1 + 2.760842} = 0.2659.$$

Question #160**Key: E**

The cdf is $F(x) = \int_0^x 4(1+t)^{-5} dt = -(1+t)^{-4} \Big|_0^x = 1 - \frac{1}{(1+x)^4}$.

Observation (x)	F(x)	compare to:	Maximum difference
0.1	0.317	0, 0.2	0.317
0.2	0.518	0.2, 0.4	0.318
0.5	0.802	0.4, 0.6	0.402
0.7	0.880	0.6, 0.8	0.280
1.3	0.964	0.8, 1.0	0.164

K-S statistic is 0.402.

Question #161**Key: D**

This follows from the formula $MSE(\hat{\theta}) = Var(\hat{\theta}) + [bias(\hat{\theta})]^2$. If the bias is zero, then the mean-squared error is equal to the variance.

Question #162**Key: B**

$E[x-d | x > d]$ is the expected payment per payment with an ordinary deductible of d . It can be evaluated (for Pareto) as

$$\begin{aligned} \frac{E(x) - E(x \wedge d)}{1 - F(d)} &= \frac{\frac{\theta}{\alpha-1} - \frac{\theta}{\alpha-1} \left[1 - \left(\frac{\theta}{d+\theta} \right)^{\alpha-1} \right]}{1 - \left[1 - \left(\frac{\theta}{d+\theta} \right)^{\alpha} \right]} \\ &= \frac{\frac{\theta}{\alpha-1} \left(\frac{\theta}{d+\theta} \right)^{\alpha-1}}{\left(\frac{\theta}{d+\theta} \right)^{\alpha}} \\ &= \frac{d+\theta}{\alpha-1} \\ &= d+\theta \text{ in this problem, since } \alpha = 2 \end{aligned}$$

$$E[x-100|x > 100] = \frac{5}{3} E[x-50|x > 50]$$

$$100 + \theta = \frac{5}{3}(50 + \theta)$$

$$300 + 3\theta = 250 + 5\theta$$

$$= \theta = 25$$

$$E[x-150|x > 150] = 150 + \theta$$

$$= 150 + 25$$

$$= 175$$

Question #163

Key: D

Let S = score

$$E(S) = E(E(S|\theta)) = E(\theta) = 75$$

$$\text{Var}(S) = E[\text{Var}(S|\theta)] + \text{Var}[E(S|\theta)]$$

$$= E(8^2) + \text{Var}(\theta)$$

$$= 64 + 6^2$$

$$= 100$$

S is normally distributed (a normal mixture of normal distributions with constant variance is normal; see Example 4.30 in Loss Models for the specific case, as we have here, with a normally distributed mean and constant variance)

$$\begin{aligned} \text{Prob}[S < 90 | S > 65] &= \frac{F(90) - F(65)}{1 - F(65)} \\ &= \frac{\Phi\left(\frac{90-75}{10}\right) - \Phi\left(\frac{65-75}{10}\right)}{1 - \Phi\left(\frac{65-75}{10}\right)} \end{aligned}$$

$$\frac{\Phi(1.5) - \Phi(-1.0)}{1 - \Phi(-1.0)} = \frac{0.9332 - (1 - 0.8413)}{1 - (1 - 0.8413)} = \frac{0.7745}{0.8413} = 0.9206$$

Note that (though this insight is unnecessary here), this is equivalent to per payment model with a franchise deductible of 65.

Question #164**Key: B**

(Referring to the number of losses, X , was a mistake. X is the random variable for the loss amount, the severity distribution).

Losses in excess of the deductible occur at a Poisson rate
of $\lambda^* = (1 - F(30))\lambda = 0.75 \times 20 = 15$

$$E(X - 30 | X > 30) = \frac{70 - 25}{0.75} = \frac{45}{0.75} = 60$$

$$\begin{aligned} \text{Var}(S) &= \lambda^* \times E((X - 30)^2 | X > 30) \\ &= 15E(X^2 - 60X + 900 | X > 30) = 15E(X^2 - 60(X - 30) - 900 | X > 30) \\ &= 15(9,000 - 60 \times 60 - 900) \\ &= 67,500 \end{aligned}$$

Question #165**Key: A**

S	$(S - 3)_+$	$E[(S - 3)_+] = E[S] - 3 + 3f_S(0) + 2f_S(1) + 1f_S(2)$
0	0	$E[S] = 2 \times [0.6 + 2 \times 0.4] = 2.8$
1	0	$f_S(0) = e^{-2}$
2	0	$f_S(1) = e^{-2} \times 2 \times (0.6) = 1.2e^{-2}$
3	0	$f_S(2) = e^{-2} \times 2(0.4) + \frac{e^{-2} 2^2}{2} \times (0.6)^2 = 1.52e^{-2}$
4	1	
5	2	
6	3	
\vdots	\vdots	

$$\begin{aligned} E[(S - 3)_+] &= 2.8 - 3 + 3 \times e^{-2} + 2 \times 1.2e^{-2} + 1 \times 1.52e^{-2} \\ &= -0.2 + 6.92e^{-2} \\ &= 0.7365 \end{aligned}$$

Question #166**Key: C**

Write (i) as $\frac{p_k}{p_{k-1}} = c + \frac{c}{k}$

This is an (a, b, 0) distribution with $a = b = c$.

Which?

1. If Poisson, $a = 0$, so $c = 0$ and $b = 0$
 $p_1 = p_2 = \dots = 0$
 $p_0 = 0.5$
 p_k 's do not sum to 1. Impossible. Thus not Poisson
2. If Geometric, $b = 0$, so $c = 0$ and $a = 0$
 By same reasoning as #1, impossible, so not Geometric.
3. If binomial, a and b have opposite signs. But here $a = b$, so not binomial.
4. Thus negative binomial.

$$1 = \frac{a}{b} = \frac{\beta/(1+\beta)}{(r-1)\beta/(1-\beta)} = \frac{1}{r-1}$$

so $r = 2$

$$p_0 = 0.5 = (1+\beta)^{-r} = (1+\beta)^{-2}$$

$$1+\beta = \sqrt{2} = 1.414$$

$$\beta = \sqrt{2} - 1 = 0.414$$

$$c = a = \beta/(1+\beta) = 0.29$$

Question #167**Key: B**

The variance calculation assumes independence, which should have been explicitly stated.

$$E(S) = E(N)E(X)$$

$$\text{Var}(S) = E(N)\text{Var}(X) + E^2(X)\text{Var}(N)$$

	$E(N)$	$\text{Var}(N)$	$E(X)$	$\text{Var}(X)$	$E(S)$	$\text{Var}(S)$
P.B	30	21	300	10,000	9,000	2.19×10^6
S.B	30	27	1000	400,000	30,000	39×10^6
L.Y	30	12	5000	2,000,000	150,000	360×10^6
					189,000	400×10^6
						(rounded)

$$\text{Standard deviation} = \sqrt{400 \times 10^6} = 20,000$$

$$189,000 + 20,000 = 209,000$$

Question #168**Key: B**

$$S_X(150) = 1 - 0.2 = 0.8$$

$$f_{Y^p}(y) = \frac{f_X(y+150)}{S_X(150)} \quad \text{So } f_{Y^p}(50) = \frac{0.2}{0.8} = 0.25$$

$$f_{Y^p}(150) = \frac{0.6}{0.8} = 0.75$$

$$E(Y^p) = (0.25)(50) + (0.75)(150) = 125$$

$$E\left[(Y^p)^2\right] = (0.25)(50^2) + (0.75)(150)^2 = 17,500$$

$$\text{Var}(Y^p) = E\left[(Y^p)^2\right] - [E(Y^p)]^2 = 17,500 - 125^2 = 1875$$

Slight time saver, if you happened to recognize it:

$$\text{Var}(Y^p) = \text{Var}(Y^p - 50) \quad \text{since subtracting a constant does not change variance, regardless of the distribution}$$

But $Y^p - 50$ takes on values only 0 and 100, so it can be expressed as 100 times a binomial random variable with $n = 1$, $q = 0.75$

$$\text{Var} = (100^2)(1)(0.25)(0.75) = 1875$$

Question #169**Key: A**

Model Solution:

X denotes the loss variable.

X_1 denotes Pareto with $\alpha = 2$; X_2 denotes Pareto with $\alpha = 4$

$$F_X(200) = 0.8 F_{X_1}(200) + 0.2 F_{X_2}(200)$$

$$= 0.8 \left[1 - \left(\frac{100}{200+100} \right)^2 \right] + 0.2 \left[1 - \left(\frac{3000}{3000+200} \right)^4 \right]$$

$$= 1 - 0.8 \left(\frac{1}{3} \right)^2 - 0.2 \left(\frac{15}{16} \right)^4$$

$$= 0.7566$$

Question #170**Key: B**

Let c denote child; ANS denote Adult Non-Smoker; AS denote Adult Smoker.

$$P(3|c)P(c) = \frac{3^3 e^{-3}}{3!} \times 0.3 = 0.067$$

$$P(3|ANS)P(ANS) = \frac{1e^{-1}}{3!} \times 0.6 = 0.037$$

$$P(3|AS)P(AS) = \frac{4^3 e^{-4}}{3!} \times 0.1 = 0.020$$

$$P(AS|N=3) = \frac{0.020}{(0.067 + 0.037 + 0.020)} = 0.16$$

Question #171**Key: C**

$$E[S] = E[N]E[X] = 3 \times 10 = 30$$

$$\begin{aligned} \text{Var}(S) &= E[N]\text{Var}(X) + E[X]^2 \text{Var}(N) \\ &= 3 \times \frac{400}{12} + 100 \times 3.6 = 100 + 360 = 460 \end{aligned}$$

For 95th percentile, $E[S] + 1.645\sqrt{\text{Var}(S)} = 30 + \sqrt{460} \times 1.645 = 65.28$

Question #172**Key: D**

The CDF is $F(x) = 1 - \frac{1}{(1+x)^4}$

Observation (x)	F(x)	compare to:	Maximum difference
0.2	0.518	0, 0.2	0.518
0.7	0.880	0.2, 0.4	0.680
0.9	0.923	0.4, 0.6	0.523
1.1	0.949	0.6, 0.8	0.349
1.3	0.964	0.8, 1.0	0.164

The K-S statistic is the maximum from the last column, 0.680.

Critical values are: 0.546, 0.608, 0.662, and 0.729 for the given levels of significance. The test statistic is between 0.662 (2.5%) and 0.729 (1.0%) and therefore the test is rejected at 0.025 and not at 0.01.

Question #173

Key: E

For claim severity,

$$\mu_S = 1(0.4) + 10(0.4) + 100(0.2) = 24.4,$$

$$\sigma_S^2 = 1^2(0.4) + 10^2(0.4) + 100^2(0.2) - 24.4^2 = 1,445.04.$$

For claim frequency,

$$\mu_F = r\beta = 3r, \quad \sigma_F^2 = r\beta(1 + \beta) = 12r.$$

For aggregate losses,

$$\mu = \mu_S \mu_F = 24.4(3r) = 73.2r,$$

$$\sigma^2 = \mu_S^2 \sigma_F^2 + \sigma_S^2 \mu_F = 24.4^2(12r) + 1,445.04(3r) = 11,479.44r.$$

For the given probability and tolerance, $\lambda_0 = (1.96/0.1)^2 = 384.16$.

The number of observations needed is

$$\lambda_0 \sigma^2 / \mu^2 = 384.16(11,479.44r) / (73.2r)^2 = 823.02 / r.$$

The average observation produces $3r$ claims and so the required number of claims is $(823.02 / r)(3r) = 2,469$.

Question #174

Key: A

$$\hat{H}(t_2) = \frac{1}{n} + \frac{1}{n-1} = \frac{2n-1}{n(n-1)} = \frac{39}{380} \Rightarrow 39n^2 - 799n + 380 = 0 \Rightarrow n = 20, n = 0.487.$$

Discard the non-integer solution to have $n = 20$.

The Kaplan-Meier Product-Limit Estimate is:

$$\hat{S}(t_9) = \frac{19}{20} \frac{18}{19} \dots \frac{11}{12} \frac{11}{20} = 0.55.$$

Question #175

Key: E

There are 27 possible bootstrap samples, which produce four different results. The results, their probabilities, and the values of g are:

Bootstrap

Sample	Prob	g
1, 1, 1	8/27	0
1, 1, 4	12/27	2
1, 4, 4	6/27	-2
4, 4, 4	1/27	0

The third central moment of the original sample is 2. Then,

$$\text{MSE} = \left[\frac{8}{27}(0-2)^2 + \frac{12}{27}(2-2)^2 + \frac{6}{27}(-2-2)^2 + \frac{1}{27}(0-2)^2 \right] = \frac{44}{9}.$$

Question #176

Key: A

Pick one of the points, say the fifth one. The vertical coordinate is $F(30)$ from the model and should be slightly less than 0.6. Inserting 30 into the five answers produces 0.573, 0.096, 0.293, 0.950, and something less than 0.5. Only the model in answer A is close.

Question #177

Key: C, E

The distribution of Θ is Pareto with parameters 1 and 2.6. Then,

$$v = EPPV = E(\Theta) = \frac{1}{2.6-1} = 0.625, \quad a = VHM = Var(\Theta) = \frac{2}{1.6(0.6)} - 0.625^2 = 1.6927,$$

$$k = v/a = 0.625/1.6927 = 0.3692, \quad Z = \frac{5}{5+0.3692} = 0.9312.$$

Question #178

DELETED

Question #179

Key: D

For an exponential distribution, the maximum likelihood estimate of θ is the sample mean, 6. Let $Y = X_1 + X_2$ where each X has an exponential distribution with mean 6.

The sample mean is $Y/2$ and Y has a gamma distribution with parameters 2 and 6.

$$\Pr(Y/2 > 10) = \Pr(Y > 20) = \int_{20}^{\infty} \frac{xe^{-x/6}}{36} dx$$

Then

$$= -\frac{xe^{-x/6}}{6} - e^{-x/6} \Big|_{20}^{\infty} = \frac{20e^{-20/6}}{6} + e^{-20/6} = 0.1546.$$

Question #180**Key: A**

From Question 9, $F(10) = 1 - \frac{10e^{-10/\theta}}{\theta} - e^{-10/\theta} = 1 - e^{-10/\theta}(1 + 10\theta^{-1}) = g(\theta)$.

$$g'(\theta) = -\frac{20}{\theta^2}e^{-20/\theta}(1 + 20\theta^{-1}) + e^{-20/\theta} \frac{20}{\theta^2} = -\frac{400e^{-20/\theta}}{\theta^3}.$$

At the maximum likelihood estimate of 6, $g'(6) = -0.066063$.

The maximum likelihood estimator is the sample mean. Its variance is the variance of one observation divided by the sample size. For the exponential distribution the variance is the square of the mean, so the estimated variance of the sample mean is $36/2 = 18$. The answer is $(-0.066063)^2(18) = 0.079$.

Question #181**Key B**

$$\mu(\lambda, \theta) = E(S | \lambda, \theta) = \lambda\theta,$$

$$v(\lambda, \theta) = \text{Var}(S | \lambda, \theta) = \lambda 2\theta^2,$$

$$v = \text{EVPV} = E(\lambda 2\theta^2) = 1(2)(1+1) = 4,$$

$$a = \text{VHM} = \text{Var}(\lambda\theta) = E(\lambda^2)E(\theta^2) - [E(\lambda)E(\theta)]^2 = 2(2) - 1 = 3,$$

$$k = v/a = 4/3.$$

Question #182**Key: B**

The distribution is binomial with $m = 100$ and $q = 0.03$. The first three probabilities are:

$$p_0 = 0.97^{100} = 0.04755, \quad p_1 = 100(0.97)^{99}(0.03) = 0.14707,$$

$$p_2 = \frac{100(99)}{2}(0.97)^{98}(0.03)^2 = 0.22515.$$

Values between 0 and 0.04755 simulate a 0, between 0.04755 and 0.19462 simulate a 1, and between 0.19462 and 0.41977 simulate a 2. The three simulated values are 2, 0, and 1. The mean is 1.

Question #183**Key: A**

A mixture of two Poissons or negative binomials will always have a variance greater than its mean. A mixture of two binomials could have a variance equal to its mean, because a single binomial has a variance less than its mean.

Question #184**Key: D**

The posterior distribution can be found from

$$\pi(\lambda | 10) \propto \frac{e^{-\lambda} \lambda^{10}}{10!} \left(\frac{0.4}{6} e^{-\lambda/6} + \frac{0.6}{12} e^{-\lambda/12} \right) \propto \lambda^{10} (0.8e^{-7\lambda/6} + 0.6e^{-13\lambda/12}).$$

The required constant is found from

$$\int_0^{\infty} \lambda^{10} (0.8e^{-7\lambda/6} + 0.6e^{-13\lambda/12}) d\lambda = 0.8(10!)(6/7)^{11} + 0.6(10!)(12/13)^{11} = 0.395536(10!).$$

The posterior mean is

$$\begin{aligned} E(\lambda | 10) &= \frac{1}{0.395536(10!)} \int_0^{\infty} \lambda^{11} (0.8e^{-7\lambda/6} + 0.6e^{-13\lambda/12}) d\lambda \\ &= \frac{0.8(11!)(6/7)^{12} + 0.6(11!)(12/13)^{12}}{0.395536(10!)} = 9.88. \end{aligned}$$

Question #185**Key: A**

$$\hat{H}(4.5) = \frac{2}{12} + \frac{1}{10} + \frac{2}{9} + \frac{2}{7} = 0.77460.$$

The variance estimate is

$$\frac{2}{12^2} + \frac{1}{10^2} + \frac{2}{9^2} + \frac{2}{7^2} = 0.089397.$$

The confidence interval is

$$0.77460 \pm 1.96\sqrt{0.089397} = 0.77460 \pm 0.58603. \text{ The interval is from 0.1886 to 1.3606.}$$

Question #186**Key: D**

$$\text{bias} = E(\hat{\theta}) - \theta = \frac{k}{k+1}\theta - \theta = -\frac{\theta}{k+1},$$

$$\text{Var}(\hat{\theta}) = \text{Var}\left(\frac{k\theta}{k+1}\right) = \frac{k^2\theta^2}{25(k+1)^2},$$

$$\text{MSE} = \text{Var}(\hat{\theta}) + \text{bias}^2 = \frac{k^2\theta^2}{25(k+1)^2} + \frac{\theta^2}{(k+1)^2},$$

$$\text{MSE} = 2\text{bias}^2 = \frac{2\theta^2}{(k+1)^2}.$$

Setting the last two equal and canceling the common terms gives

$$\frac{k^2}{25} + 1 = 2 \text{ for } k = 5.$$

Question #187

Key: D

For the geometric distribution $\mu(\beta) = \beta$ and $v(\beta) = \beta(1 + \beta)$. The prior density is Pareto with parameters α and 1. Then,

$$\mu = E(\beta) = \frac{1}{\alpha - 1},$$

$$v = \text{EVPV} = E[\beta(1 + \beta)] = \frac{1}{\alpha - 1} + \frac{2}{(\alpha - 1)(\alpha - 2)} = \frac{\alpha}{(\alpha - 1)(\alpha - 2)},$$

$$a = \text{VHM} = \text{Var}(\beta) = \frac{2}{(\alpha - 1)(\alpha - 2)} - \frac{1}{(\alpha - 1)^2} = \frac{\alpha}{(\alpha - 1)^2(\alpha - 2)},$$

$$k = v/a = \alpha - 1, \quad Z = \frac{1}{1+k} = \frac{1}{\alpha}.$$

The estimate is

$$\frac{1}{\alpha}x + \left(1 - \frac{1}{\alpha}\right)\frac{1}{\alpha - 1} = \frac{x+1}{\alpha}.$$

Question #188

DELETED

Question #189

Key: E

A is false. Using sample data gives a better than expected fit and therefore a test statistic that favors the null hypothesis, thus increasing the Type II error probability. The K-S test works only on individual data and so B is false. The A-D test emphasizes the

tails, thus C is false. D is false because the critical value depends on the degrees of freedom which in turn depends on the number of cells, not the sample size.

Question #190

Key: B

$$E(\theta) = 0.05(0.8) + 0.3(0.2) = 0.1,$$

$$E(\theta^2) = 0.05^2(0.8) + 0.3^2(0.2) = 0.02,$$

$$\mu(\theta) = 0(2\theta) + 1(\theta) + 2(1 - 3\theta) = 2 - 5\theta,$$

$$v(\theta) = 0^2(2\theta) + 1^2(\theta) + 2^2(1 - 3\theta) - (2 - 5\theta)^2 = 9\theta - 25\theta^2,$$

$$\mu = E(2 - 5\theta) = 2 - 5(0.1) = 1.5,$$

$$v = EVPV = E(9\theta - 25\theta^2) = 9(0.1) - 25(0.02) = 0.4,$$

$$a = VHM = Var(2 - 5\theta) = 25Var(\theta) = 25(0.02 - 0.1^2) = 0.25,$$

$$k = v/a = 0.4/0.25 = 1.6, Z = \frac{1}{1+1.6} = \frac{5}{13},$$

$$P = \frac{5}{13}2 + \frac{8}{13}1.5 = 1.6923.$$

Question #191

Key: B

$f(\lambda | 5, 3) \propto \frac{e^{-\lambda} \lambda^5}{5!} \frac{e^{-\lambda} \lambda^3}{3!} \frac{2^5 \lambda^5 e^{-2\lambda}}{24\lambda} \propto \lambda^{12} e^{-4\lambda}$. This is a gamma distribution with parameters 13 and 0.25. The expected value is $13(0.25) = 3.25$.

Alternatively, if the Poisson-gamma relationships are known, begin with the prior parameters $\alpha = 5$ and $\beta = 2$ where $\beta = 1/\theta$ if the parameterization from *Loss Models* is considered. Then the posterior parameters are $\alpha' = 5 + 5 + 3 = 13$ where the second 5 and the 3 are the observations and $\beta' = 2 + 2 = 4$ where the second 2 is the number of observations. The posterior mean is then $13/4 = 3.25$.

Question #192

Key: D

Because the kernel extends one unit each direction there is no overlap. The result will be three replications of the kernel. The middle one will be twice as high due to having two observations of 3 while only one observation at 1 and 5. Only graphs A and D fit this description. The kernel function is smooth, which rules out graph A.

Question #193**Key: C**

The two moment equations are

$$508 = \frac{\theta}{\alpha - 1}, \quad 701,401.6 = \frac{2\theta^2}{(\alpha - 1)(\alpha - 2)}. \quad \text{Dividing the square of the first equation into the}$$

$$\text{second equation gives } \frac{701,401.6}{508^2} = 2.7179366 = \frac{2(\alpha - 1)}{\alpha - 1}. \quad \text{The solution is } \alpha = 4.785761.$$

From the first equation, $\theta = 1,923.167$. The requested LEV is

$$E(X \wedge 500) = \frac{1,923.167}{3.785761} \left[1 - \left(\frac{1,923.167}{1,923.167 + 500} \right)^{3.785761} \right] = 296.21.$$

Question #194**Key: B**

$$\hat{v} = \widehat{EVPV} = \frac{50(200 - 227.27)^2 + 60(250 - 227.27)^2 + 100(160 - 178.95)^2 + 90(200 - 178.95)^2}{1 + 1}$$

$$= 71,985.647,$$

$$\hat{k} = 71,985.647 / 651.03 = 110.57,$$

$$\hat{Z} = \frac{110}{110 + 110.57} = 0.499.$$

Question #195**Key: B**

$$F_{100}(1,000) = 0.16, \quad F_{100}(3,000) = 0.38, \quad F_{100}(5,000) = 0.63, \quad F_{100}(10,000) = 0.81,$$

$$F_{100}(2,000) = 0.5(0.16) + 0.5(0.38) = 0.27,$$

$$F_{100}(6,000) = 0.8(0.63) + 0.2(0.81) = 0.666.$$

$$\Pr(2,000 < X < 6,000) = 0.666 - 0.27 = 0.396.$$

Question #196**Key: C**

$$\begin{aligned}
L &= \left[\frac{f(750)}{1-F(200)} \right]^3 f(200)^3 f(300)^4 [1-F(10,000)]^6 \left[\frac{f(400)}{1-F(300)} \right]^4 \\
&= \left[\frac{\alpha 10,200^\alpha}{10,750^{\alpha+1}} \right]^3 \left[\frac{\alpha 10,000^\alpha}{10,200^{\alpha+1}} \right]^3 \left[\frac{\alpha 10,000^\alpha}{10,300^{\alpha+1}} \right]^4 \left[\frac{10,000^\alpha}{20,000^\alpha} \right]^6 \left[\frac{\alpha 10,300^\alpha}{10,400^{\alpha+1}} \right]^4 \\
&= \alpha^{14} 10,200^{-3} 10,000^{13} \alpha 10,300^{-4} 10,750^{-3\alpha-3} 20,000^{-6\alpha} 10,400^{-4\alpha-4} \\
&\propto \alpha^{14} 10,000^{13} \alpha 10,750^{-3\alpha} 20,000^{-6\alpha} 10,400^{-4\alpha}, \\
\ln L &= 14 \ln \alpha + 13 \alpha \ln(10,000) - 3 \alpha \ln(10,750) - 6 \alpha \ln(20,000) - 4 \alpha \ln(10,400) \\
&= 14 \ln \alpha - 4.5327 \alpha.
\end{aligned}$$

The derivative is $14/\alpha - 4.5327$ and setting it equal to zero gives $\hat{\alpha} = 3.089$.

Question #197

Key: C

$$\hat{v} = EVPV = \bar{x} = \frac{30+30+12+4}{100} = 0.76.$$

$$\begin{aligned}
\hat{a} = VHM &= \frac{50(0-0.76)^2 + 30(1-0.76)^2 + 15(2-0.76)^2 + 4(3-0.76)^2 + 1(4-0.76)^2}{99} - 0.76 \\
&= 0.090909,
\end{aligned}$$

$$\hat{k} = \frac{0.76}{0.090909} = 8.36, \quad \hat{Z} = \frac{1}{1+8.36} = 0.10684,$$

$$P = 0.10684(1) + 0.89316(0.76) = 0.78564.$$

The above analysis was based on the distribution of total claims for two years. Thus 0.78564 is the expected number of claims for the next two years. For the next one year the expected number is $0.78564/2 = 0.39282$.

Question #198

DELETED

Question #199**Key: E**

The density function is $f(x) = \frac{0.2x^{-0.8}}{\theta^{0.2}} e^{-(x/\theta)^{0.2}}$. The likelihood function is

$$L(\theta) = f(130)f(240)f(300)f(540)[1 - F(1000)]^2$$

$$= \frac{0.2(130)^{-0.8}}{\theta^{0.2}} e^{-(130/\theta)^{0.2}} \frac{0.2(240)^{-0.8}}{\theta^{0.2}} e^{-(240/\theta)^{0.2}} \frac{0.2(300)^{-0.8}}{\theta^{0.2}} e^{-(300/\theta)^{0.2}} \frac{0.2(540)^{-0.8}}{\theta^{0.2}} e^{-(540/\theta)^{0.2}} e^{-(1000/\theta)^{0.2}} e^{-(1000/\theta)^{0.2}}$$

$$\propto \theta^{-0.8} e^{-\theta^{-0.2}(130^{0.2} + 240^{0.2} + 300^{0.2} + 540^{0.2} + 1000^{0.2} + 1000^{0.2})}$$

$$l(\theta) = -0.8 \ln(\theta) - \theta^{-0.2}(130^{0.2} + 240^{0.2} + 300^{0.2} + 540^{0.2} + 1000^{0.2} + 1000^{0.2})$$

$$= -0.8 \ln(\theta) - 20.2505\theta^{-0.2},$$

$$l'(\theta) = -0.8\theta^{-1} + 0.2(20.2505)\theta^{-1.2} = 0,$$

$$\theta^{-0.2} = 0.197526, \quad \hat{\theta} = 3,325.67.$$

Question #200**Key: A**

Buhlmann estimates are on a straight line, which eliminates E. Bayes estimates are never outside the range of the prior distribution. Because graphs B and D include values outside the range 1 to 4, they cannot be correct. The Buhlmann estimates are the linear least squares approximation to the Bayes estimates. In graph C the Bayes estimates are consistently higher and so the Buhlmann estimates are not the best approximation. This leaves A as the only feasible choice.

Question #201**Key: C**

The expected counts are $300(0.035) = 10.5$, $300(0.095) = 28.5$, $300(0.5) = 150$, $300(0.2) = 60$, and $300(0.17) = 51$ for the five groups. The test statistic is

$$\frac{(5-10.5)^2}{10.5} + \frac{(42-28.5)^2}{28.5} + \frac{(137-150)^2}{150} + \frac{(66-60)^2}{60} + \frac{(50-51)^2}{51} = 11.02.$$

There are $5 - 1 = 4$ degrees of freedom. From the table, the critical value for a 5% test is 9.488 and for a 2.5% test is 11.143. The hypothesis is rejected at 5%, but not at 2.5%.

Question #202**Key: A**

To simulate a lognormal variate, first convert the uniform random number to a standard normal. This can be done by using the table of normal distribution values. For the six values given, the corresponding standard normal values are 0.3, -0.1, 1.6, -1.4, 0.8, and -0.2. Next, multiply each number by the standard deviation of 0.75 and add the mean of

5.6. This produces random observations from the normal 5.6, 0.75² distribution. These values are 5.825, 5.525, 6.8, 4.55, 6.2, and 5.45. To create lognormal observations, exponentiate these values. The results are 339, 251, 898, 95, 493, and 233. After imposing the policy limit of 400, the mean is (339 + 251 + 400 + 95 + 400 + 233)/6 = 286.

Question #203

Key: C

For the geometric distribution, $\Pr(X_1 = 2 | \beta) = \frac{\beta^2}{(1 + \beta)^3}$ and the expected value is β .

$$\Pr(\beta = 2 | X_1 = 2) = \frac{\Pr(X_1 = 2 | \beta = 2) \Pr(\beta = 2)}{\Pr(X_1 = 2 | \beta = 2) \Pr(\beta = 2) + \Pr(X_1 = 2 | \beta = 5) \Pr(\beta = 5)}$$

$$= \frac{\frac{4}{27} \frac{1}{3}}{\frac{4}{27} \frac{1}{3} + \frac{25}{216} \frac{1}{3}} = 0.39024.$$

The expected value is then $0.39024(2) + 0.60976(5) = 3.83$.

Question #204

Key: D

The following derives the general formula for the statistic to be forgotten by time x. It would work fine, and the equations would look simpler, if you immediately plugged in

$x = \frac{1}{2}$, the only value you want. Then the $x + \frac{1}{2}$ becomes 1.

Let X be the random variable for when the statistic is forgotten. Then $F_X(x|y) = 1 - e^{-xy}$

For the unconditional distribution of X, integrate with respect to y

$$F_X(x) = \int_0^{\infty} (1 - e^{-xy}) \frac{1}{\Gamma(2)y} \left(\frac{y}{2}\right)^2 e^{-y/2} dy$$

$$= 1 - \frac{1}{4} \int_0^{\infty} y e^{-y(x+1/2)} dy$$

$$= 1 - \frac{1}{4(x+1/2)^2}$$

$$F(1/2) = 1 - \frac{1}{4(1/2+1/2)^2} = 0.75$$

There are various ways to evaluate the integral in the second line:

1. Calculus, integration by parts

2. Recognize that $\int_0^\infty y \left(x + \frac{1}{2}\right) e^{-y\left(x + \frac{1}{2}\right)} dy$

is the expected value of an exponential random variable with $\theta = \frac{1}{x + \frac{1}{2}}$

3. Recognize that $\Gamma(2) \left(x + \frac{1}{2}\right)^2 y e^{-y\left(x + \frac{1}{2}\right)}$ is the density function for a Gamma

random variable with $\alpha = 2$ and $\theta = \frac{1}{x + \frac{1}{2}}$, so it would integrate to 1.

(Approaches 2 and 3 would also work if you had plugged in $x = \frac{1}{2}$ at the start. The resulting θ becomes 1).

Question #205

Key: D

State#	Number	Probability of needing Therapy	Mean Number of visits E(X)	E(N)	Var(N)	Var(X)	E(S)	Var(S)
1	400	0.2	2	80	64	6	160	736
2	300	0.5	15	150	75	240	2,250	52,875
3	200	0.3	9	60	42	90	540	8,802
							2,950	62,413

$$\text{Std Dev } (S) = \sqrt{62413} = 250$$

$$\Pr(S > 3000) = \Pr\left(\frac{S - 2950}{250} > \frac{50}{250}\right) = 1 - \Phi(0.2) = 0.42$$

The $\text{Var}(X)$ column came from the formulas for mean and variance of a geometric distribution.

Using the continuity correction, solving for $\Pr(S > 3000.5)$, is theoretically better but does not affect the rounded answer.

Question #206**Key: B**Frequency is geometric with $\beta = 2$, so

$$p_0 = 1/3, \quad p_1 = 2/9, \quad p_2 = 4/27$$

Convolutions of $f_X(x)$ needed are

X	f	f^{*2}
5	0.2	0
10	0.3	0.04

$$\text{so } f_S(0) = 1/3, \quad f_S(5) = 2/9(0.2) = 0.044, \quad f_S(10) = 2/9(0.3) + 4/27(0.04) = 0.073$$

$$E(X) = (0.2)(5) + (0.3)(10) + (0.5)(20) = 14$$

$$E[S] = 2E(X) = 28$$

$$\begin{aligned} E[S-15]_+ &= E[S] - 5(1-F(0)) - 5(1-F(5)) - 5(1-F(10)) \\ &= 28 - 5(1-1/3) - 5(1-1/3-0.044) - 5(1-1/3-0.044-0.073) \\ &= 28 - 3.33 - 3.11 - 2.75 = 18.81 \end{aligned}$$

Alternatively,

$$\begin{aligned} E[S-15]_+ &= E[S] - 15 + 15f_S(0) + 10f_S(5) + 5f_S(10) \\ &= 28 - 15 + (15)\left(\frac{1}{3}\right) + 10(0.044) + 5(0.073) \\ &= 18.81 \end{aligned}$$

Question #207**Key: E**

$$\begin{aligned} S_X(4) &= 1 - \int_0^4 f_X(x) dx = 1 - \int_0^4 0.02x dx \\ &= 1 - 0.01x^2 \Big|_0^4 \\ &= 0.84 \end{aligned}$$

$$f_{Y^p}(y) = \frac{f_X(y+4)}{S_X(4)} = \frac{0.02(y+4)}{0.84} = 0.0238(y+4)^2$$

$$E(Y^p) = \int_0^6 y(0.0238(y+4))dy = 0.0238 \left(\frac{y^3}{3} + \frac{4y^2}{2} \right) \Big|_0^6$$

$$= 3.4272$$

Question #208

Key: E

By Theorem 4.51 (on page 93 of the second edition of Loss Models), probability of zero claims = pgf of negative binomial applied to the probability that Poisson equals 0.

For the Poisson, $f(0) = e^{-\lambda}$

$$\text{So } 0.067 = [1 - \beta(e^{-\lambda} - 1)]^{-r} = [1 - 3(e^{-\lambda} - 1)]^{-2}$$

Solving gives $\lambda = 3$

Question #209

Key: D

For any deductible d and the given severity distribution

$$E(X-d)_+ = E(X) - E(X \wedge d)$$

$$= 3000 - 3000 \left(1 - \frac{3000}{3000+d} \right)$$

$$= (3000) \left(\frac{3000}{3000+d} \right)$$

$$\text{So } P_{2005} = (1.2)(3000) \left(\frac{3000}{3600} \right) = 3000$$

The following paragraph just clarifies the notation in the rest of the solution:

Let r denote the reinsurer's deductible relative to losses (not relative to reinsured claims). Thus if $r = 1000$ (we are about to solve for r), then on a loss of 4000, the insured collects

$4000 - 600 = 3400$, the reinsurer pays $4000 - 1000 = 3000$, leaving the primary insurer paying 400.

Another way, exactly equivalent, to express that reinsurance is that the primary company pays the insured 3400. The reinsurer reimburses the primary company for its claims less a deductible of 400 applied to claims. So the reinsurer pays $3400 - 400 = 3000$, the same as before.

Expected reinsured claims in 2005

$$= (3000) \left(\frac{3000}{3000+r} \right) = \frac{9,000,000}{3000+r}$$

$$R_{2005} = (1.1) \left(\frac{9,000,000}{3000+r} \right) = (0.55) P_{2005}$$

$$\frac{9,900,000}{3000+r} = (0.55)(3000) = 1650$$

$$r = 3000$$

In 2006, after 20% inflation, losses will have a two-parameter Pareto distribution with $\alpha = 2$ and $\theta = (1.2)(3000) = 3600$.

The general formula for claims will be

$$E(X-d)_+ = (3600) \left(\frac{3600}{3600+d} \right) = \frac{12,960,000}{3600+d}$$

$$P_{2006} = 1.2 \left(\frac{12,960,000}{3000+600} \right) = 3703$$

$$R_{2006} = 1.1 \left(\frac{12,960,000}{3600+3000} \right) = 2160$$

$$R_{2006} / P_{2006} = 0.5833$$

[If you applied the reinsurer's deductible to the primary insurer's claims, you would solve that the deductible is 2400, and the answer to the problem is the same].

Question #210

Key: C

Consider Disease 1 and other Diseases as independent Poisson processes with respective λ 's $= (0.16) \left(\frac{1}{16} \right) = 0.01$ and $(0.16) \left(\frac{15}{16} \right) = 0.15$ respectively. Let $S_1 =$ aggregate losses from Disease 1; $S_2 =$ aggregate losses from other diseases.

$$E(S_1) = 100 \times 0.01 \times 5 = 5$$

$$\text{Var}(S_1) = 100 \times 0.01 \times (5^2 + 5^2) = 2525$$

$$E(S_2) = 100 \times 0.15 \times 10 = 150$$

$$\text{Var}(S_2) = 100 \times 0.15 \times (20^2 + 10^2) = 7500$$

If no one gets the vaccine:

$$E(S) = 5 + 150 = 155$$

$$\text{Var}(S) = 2525 + 7500 = 10,025$$

$$\Phi(0.7) = 1 - 0.24$$

$$A = 155 + 0.7\sqrt{10,025} = 225.08$$

If all get the vaccine, vaccine cost = $(100)(0.15) = 15$

No cost or variance from Disease 1

$$B = 15 + 150 + 0.7\sqrt{7500} = 225.62$$

$$A/B = 0.998$$

Question #211

Key: A

For current model $f(x) = \frac{1}{4}e^{-x/4}$

Let $g(x)$ be the new density function, which has

(i) $g(x) = c, \quad 0 \leq x \leq 3$

(ii) $g(x) = ke^{-x/4}, \quad x > 3^*$

(iii) $c = ke^{-3/4}$, since continuous at $x = 3$

Since g is density function, it must integrate to 1.

$$1 = 3c + \int_3^{\infty} ke^{-x/4} dx = 3ke^{-3/4} + 4ke^{-3/4} = 3c + 4c \Rightarrow c = \frac{1}{7}$$

$$F(3) = \int_0^3 c dx = \int_0^3 \frac{1}{7} dx = \frac{3}{7} = 0.43$$

*This could equally well have been written $g(x) = d \times \left(\frac{1}{4}e^{-x/4}\right)$, then let $k = d/4$, or even carry the $d/4$ throughout.

Question #212

Key: C

Since loss amounts are uniform on $(0, 10)$, 40% of losses are below the deductible (4), and 60% are above. Thus, claims occur at a Poisson rate $\lambda^* = (0.6)(10) = 6$.

Since loss amounts were uniform on $(0, 10)$, claims are uniform on $(0, 6)$.

Let N = number of claims; X = claim amount; S = aggregate claims.

$$E(N) = \text{Var}(N) = \lambda^* = 6$$

$$E(X) = (6-0)/2 = 3$$

$$\text{Var}(X) = (6-0)^2/12 = 3$$

$$\begin{aligned} \text{Var}(S) &= E(N)\text{Var}(X) + \text{Var}(N) [E(X)]^2 \\ &= 6*3 + 6*3^2 \\ &= 72 \end{aligned}$$

Question #213

Key: E

N	p_n	$n \times p_n$	$n^2 \times p_n$
0	0.1	0	0
1	0.4	0.4	0.4
2	0.3	0.6	1.2
3	0.2	0.6	1.8
		$E[N] = 1.6$	$E[N^2] = 3.4$

$$\text{Var}(N) = 3.4 - 1.6^2 = 0.84$$

$$E[X] = \lambda = 3$$

$$\text{Var}(X) = \lambda = 3$$

$$\begin{aligned} \text{Var}(S) &= \\ &= E[N]\text{Var}(X) + E(X)^2 \times \text{Var}(N) \\ &= 1.6(3) + (3)^2(0.84) \\ &= 12.36 \end{aligned}$$

Question #214

Key: D

$$\hat{S}(300) = 3/10 \text{ (there are three observations greater than 300)}$$

$$\hat{H}(300) = -\ln[\hat{S}(300)] = -\ln(0.3) = 1.204.$$

Question #215**Key: A**

$$E(X | \lambda) = \text{Var}(X | \lambda) = \lambda$$

$$\mu = v = E(\lambda) = \alpha\theta; a = \text{Var}(\lambda) = \alpha\theta^2; k = v/a = 1/\theta$$

$$Z = \frac{n}{n+1/\theta} = \frac{n\theta}{n\theta+1}$$

$$0.15 = \frac{\theta}{\theta+1}(1) + \frac{1}{\theta+1}\mu = \frac{\theta+\mu}{\theta+1}$$

$$0.20 = \frac{2\theta}{2\theta+1}(2) + \frac{1}{2\theta+1}\mu = \frac{4\theta+\mu}{2\theta+1}$$

From the first equation,

$$0.15\theta + 0.15 = \theta + \mu \text{ and so } \mu = 0.15 - 0.85\theta$$

Then the second equation becomes

$$0.4\theta + 0.2 = 4\theta + 0.15 - 0.85\theta$$

$$0.05 = 2.75\theta; \theta = 0.01818$$

Question #216**Key: E**

$$0.75 = \frac{1}{1+(100/\theta)^\gamma}; 0.25 = \frac{1}{1+(500/\theta)^\gamma}$$

$$(100/\theta)^\gamma = 1/3; (500/\theta)^\gamma = 3$$

Taking the ratio of these two equalities produces $5^\gamma = 9$. From the second equality,

$$9 = [(500/\theta)^2]^\gamma = 5^\gamma; (500/\theta)^2 = 5; \theta = 223.61$$

Question #217**Key: E**

Begin with

y	350	500	1000	1200	1500
s	2	2	1	1	1
r	10	8	5	2	1

$$\text{Then } \hat{S}_1(1250) = \frac{8}{10} \frac{6}{8} \frac{4}{5} \frac{1}{2} = 0.24$$

The likelihood function is

$$L(\theta) = \left[\theta^{-1} e^{-350/\theta} \right]^2 \left[\theta^{-1} e^{-500/\theta} \right]^2 e^{-500/\theta} \theta^{-1} e^{-1000/\theta} \left[e^{-1000/\theta} \right]^2 \theta^{-1} e^{-1200/\theta} \theta^{-1} e^{-1500/\theta}$$

$$= \theta^{-7} e^{-7900/\theta}$$

$$l(\theta) = -7 \ln \theta - \frac{7900}{\theta}; l'(\theta) = -\frac{7}{\theta} + \frac{7900}{\theta^2} = 0; \hat{\theta} = 7900/7$$

$$\hat{S}_2(1250) = e^{-1250(7)/7900} = 0.33$$

The absolute difference is 0.09.

Question #218**Key: E**

$$f(x) = -S'(x) = \frac{4x\theta^4}{(\theta^2 + x^2)^3}$$

$$L(\theta) = f(2)f(4)S(4) = \frac{4(2)\theta^4}{(\theta^2 + 2^2)^3} \frac{4(4)\theta^4}{(\theta^2 + 4^2)^3} \frac{\theta^4}{(\theta^2 + 4^2)^2} = \frac{128\theta^{12}}{(\theta^2 + 4)^3(\theta^2 + 16)^5}$$

$$l(\theta) = \ln 128 + 12 \ln \theta - 3 \ln(\theta^2 + 4) - 5 \ln(\theta^2 + 16)$$

$$l'(\theta) = \frac{12}{\theta} - \frac{6\theta}{\theta^2 + 4} - \frac{10\theta}{\theta^2 + 16} = 0; 12(\theta^4 + 20\theta^2 + 64) - 6(\theta^4 + 16\theta^2) - 10(\theta^4 + 4\theta^2) = 0$$

$$0 = -4\theta^4 + 104\theta^2 + 768 = \theta^4 - 26\theta^2 - 192$$

$$\theta^2 = \frac{26 \pm \sqrt{26^2 + 4(192)}}{2} = 32; \theta = 5.657$$

Question #219**Key: A**

$$E(X | \theta) = \int_0^\theta x \frac{2x}{\theta^2} dx = \frac{2\theta}{3}; \text{Var}(X | \theta) = \int_0^\theta x^2 \frac{2x}{\theta^2} dx - \frac{4\theta^2}{9} = \frac{\theta^2}{2} - \frac{4\theta^2}{9} = \frac{\theta^2}{18}$$

$$\mu = (2/3)E(\theta) = (2/3) \int_0^1 4\theta^4 d\theta = 8/15$$

$$EVPV = v = (1/18)E(\theta^2) = (1/18) \int_0^1 4\theta^5 d\theta = 1/27$$

$$VHM = a = (2/3)^2 \text{Var}(\theta) = (4/9) [4/6 - (4/5)^2] = 8/675$$

$$k = \frac{1/27}{8/675} = 25/8; Z = \frac{1}{1 + 25/8} = 8/33$$

Estimate is $(8/33)(0.1) + (25/33)(8/15) = 0.428$.


Question #220**Key: D**

From the Poisson(4) distribution the probabilities at 0, 1, and 2 are 0.0183, 0.0733, and 0.1463. The cumulative probabilities are 0.0183, 0.0916, and 0.2381. Because $0.0916 < 0.13 < 0.2381$ the simulated number of claims is 2. Claim amounts are simulated from solving


$u = 1 - e^{-x/1000}$ for $x = -1000 \ln(1 - u)$. The two simulated amounts are 51.29 and 2995.73 for a total of 3047.02

Question #221**Key: B**

It may be easiest to show this by graphing the density functions. For the first function the three components are each constant. One is of height 1/20 from 0 to 2 (representing the empirical probability of 1/10 at 1, one is height 1/20 from 1 to 3 and one is height 8/20 from 2 to 4. The following figure shows each of them and their sum, the kernel density estimator.



The triangular one is similar. For the triangle from 0 to 2, the area must be $1/10$. With a base of 2, the height is $1/10$. the same holds for the second triangle. The third has height $8/10$. When added they look as follows;



The question asks about cumulative probabilities. From 0 to 1 the first is linear and the second is quadratic, but by $x = 1$ both have accumulated 0.05 of probability. Because the cumulative distribution functions are the same at 1 and the density functions are identical from 1 to 2, the distribution functions must be identical from 1 to 2.

Question #222**Key: D and E**

For the Poisson distribution, the mean, λ , is estimated as $230/1000 = 0.23$.

# of Days	Poisson Probability	Expected # of Workers	Observed # of Workers	χ^2
0	0.794533	794.53	818	0.69
1	0.182743	182.74	153	4.84
2	0.021015	21.02	25	0.75
3 or more	0.001709	1.71	4	3.07
Total			1000	9.35

The χ^2 distribution has 2 degrees of freedom because there are four categories and the Poisson parameter is estimated (d.f. = 4 - 1 - 1 = 2).

The critical values for a chi-square test with two degrees of freedom are shown in the following table.

Significance Level	Critical Value
10%	4.61
5%	5.99
2.5%	7.38
1%	9.21

9.35 is greater than 9.21 so the null hypothesis is rejected at the 1% significance level.

Question #223**Key: D**

$$EVPV = \hat{v} = \frac{25(480 - 472.73)^2 + 30(466.67 - 472.73)^2}{2 - 1} = 2423.03 \text{ where } 480 = 12,000/25,$$

$466.67 = 14,000/30$, and $472.73 = 26,000/55$.

$$k = 2423.03 / 254 = 9.54; Z = \frac{55}{55 + 9.54} = 0.852$$

Question #224**DELETED**

Question #225**Key: C**

The quantity of interest is $P = \Pr(X \leq 5000) = \Phi\left(\frac{\ln 5000 - \mu}{\sigma}\right)$. The point estimate is

$$\Phi\left(\frac{\ln 5000 - 6.84}{1.49}\right) = \Phi(1.125) = 0.87.$$

For the delta method:

$$\frac{\partial P}{\partial \mu} = \frac{-\phi(1.125)}{1.49} = -0.1422; \quad \frac{\partial P}{\partial \sigma} = \frac{-1.125\phi(1.125)}{1.49} = -0.1600 \quad \text{where } \phi(z) = \frac{1}{\sqrt{2\pi}} e^{-z^2/2}.$$

Then the variance of \hat{P} is estimated as $(-0.1422)^2 0.0444 + (-0.16)^2 0.0222 = 0.001466$ and the lower limit is $P_L = 0.87 - 1.96\sqrt{0.001466} = 0.79496$.

Question #226**Key: A**

$$\begin{aligned} \Pr(\theta = 0.1 | X_1 = 1) &= \frac{\Pr(X_1 = 1 | \theta = 0.1) \Pr(\theta = 0.1)}{\Pr(X_1 = 1 | \theta = 0.1) \Pr(\theta = 0.1) + \Pr(X_1 = 1 | \theta = 0.3) \Pr(\theta = 0.3)} \\ &= \frac{0.1(0.8)}{0.1(0.8) + 0.3(0.2)} = \frac{4}{7} \end{aligned}$$

Then,

$$E(X_2 | \theta = 0.1) = 0(0.2) + 1(0.1) + 2(0.7) = 1.5$$

$$E(X_2 | \theta = 0.3) = 0(0.6) + 1(0.3) + 2(0.1) = 0.5$$

$$E(X_2 | X_1 = 1) = (1.5) \frac{4}{7} + (0.5) \frac{3}{7} = 1.071$$

Question #227**Key: D**

The requirement is that

$$0.01 \hat{F}(1500) \geq 1.96 \sqrt{\frac{\hat{F}(1500) \hat{S}(1500)}{N}}$$

$$0.0001 \frac{P^2}{N^2} \geq 3.8416 \frac{P(N-P)}{N^3}$$

$$\frac{NP}{N-P} \geq 38,416.$$

For the five answer choices, the left hand side is 34,364, 15,000, 27,125, 39,243, and 37,688. Only answer D meets the condition.

Question #228

Key: D

$$\frac{s_4}{r_4} = \hat{H}(y_4) - \hat{H}(y_3) = 0.5691 - 0.4128 = 0.1563.$$

$$\frac{s_4^2}{r_4^2} = \hat{V}[\hat{H}(y_4)] - \hat{V}[\hat{H}(y_3)] = 0.014448 - 0.009565 = 0.004883.$$

$$\text{Therefore, } s_4 = \frac{(s_4/r_4)^2}{s_4/r_4^2} = \frac{0.1563^2}{0.004833} = 5.$$

Question #229

Key: A

$$\ln f(x) = \ln \theta - 2 \ln(\theta + x)$$

$$\frac{\partial \ln f(x)}{\partial \theta} = \frac{1}{\theta} - \frac{2}{\theta + x}$$

$$\frac{\partial^2 \ln f(x)}{\partial \theta^2} = -\frac{1}{\theta^2} + \frac{2}{(\theta + x)^2}$$

$$E \left[\frac{\partial^2 \ln f(x)}{\partial \theta^2} \right] = -\frac{1}{\theta^2} + \int_0^\infty \frac{2\theta}{(\theta + x)^4} dx = -\frac{1}{\theta^2} + \left[-\frac{2\theta}{3(\theta + x)^3} \right]_0^\infty = -\frac{1}{\theta^2} + \frac{2}{3\theta^2} = -\frac{1}{3\theta^2}$$

$$I(\theta) = \frac{n}{3\theta^2}; \quad \text{Var} = \frac{3\theta^2}{n}$$

Question #230

Key: B

$$\mu = E[E(X | \lambda)] = E(\lambda) = 1(0.9) + 10(0.09) + 20(0.01) = 2$$

$$EVPV = v = E[\text{Var}(X | \lambda)] = E(\lambda) = 2$$

$$VHM = a = \text{Var}[E(X | \lambda)] = \text{Var}(\lambda) = 1(0.9) + 100(0.09) + 400(0.01) - 2^2 = 9.9$$

$$Z = \frac{1}{1 + 2/9.9} = 0.83193; \quad 11.983 = 0.83193x + 0.16807(2); \quad x = 14$$

Question #231

Key: A

The given interval for H can be written as $0.775 \pm 1.96\sqrt{0.063}$ and therefore the estimated variance of \hat{H} is 0.063. To apply the delta method,

$$S = e^{-H}; \quad \frac{dS}{dH} = -e^{-H}; \quad \text{Var}(\hat{S}) \doteq (-e^{-\hat{H}})^2 \text{Var}(\hat{H}) = (-e^{-0.775})^2 (0.063) = 0.134.$$

The point estimate of S is $e^{-0.775} = 0.4607$ and the confidence interval is $0.4607 \pm 1.96\sqrt{0.134} = 0.2269$ or $(0.23, 0.69)$.

Question #232

Key: B

The first step is to trend the year 1 data by 1.21 and the year 2 data by 1.1. The observations are now 24.2, 48.4, 60.5, 33, 44, 99, and 132.

The first two sample moments are 63.014 and 5262.64. The equations to solve are

$$63.014 = e^{\mu+0.5\sigma^2}; \quad 4.14336 = \mu + 0.5\sigma^2$$

$$5262.64 = e^{2\mu+2\sigma^2}; \quad 8.56839 = 2\mu + 2\sigma^2.$$

Taking four times the first equation and subtracting the second gives 2μ and therefore

$$\mu = \frac{4(4.14336) - 8.56839}{2} = 4.00.$$

Question #233

Key: A

$$\hat{\mu} = \bar{x} = 12 / 60 = 0.2, \quad \text{EVPV} = \hat{\nu} = \bar{x} = 0.2$$

$$\text{VHM} = \hat{a} = \frac{10(0.4 - 0.2)^2 + 20(0.25 - 0.2)^2 + 30(0.1 - 0.2)^2 - (3-1)(0.2)}{60 - \frac{10^2 + 20^2 + 30^2}{60}} = 0.009545$$

$$\hat{k} = 20.9524; \quad Z = \frac{10}{10 + 20.9524} = 0.323$$

Question #234

DELETED

Question #235

Key: C

$$l(\tau, \theta) = \sum_{j=1}^5 \ln f(x_j) = \sum_{j=1}^5 \ln \tau + (\tau - 1) \ln x_j - \tau \ln \theta - (x_j / \theta)^\tau. \quad \text{Under the null hypothesis it is}$$

$$l(2, \theta) = \sum_{j=1}^5 \ln 2 + \ln x_j - 2 \ln \theta - (x_j / \theta)^2. \quad \text{Inserting the maximizing value of 816.7 for } \theta$$

gives -35.28 . The likelihood ratio test statistic is $2(-33.05 + 35.28) = 4.46$. There is one degree of freedom. At a 5% significance level the critical value is 3.84 and at a 2.5% significance level it is 5.02.

Question #236

Key: C

It is given that $n = 4$, $v = 8$, and $Z = 0.4$. Then, $0.4 = \frac{4}{4 + \frac{8}{a}}$ which solves for $a = 4/3$. For

the covariance,

$$\begin{aligned} \text{Cov}(X_i, X_j) &= E(X_i X_j) - E(X_i)E(X_j) \\ &= E[E(X_i X_j | \theta)] - E[E(X_i | \theta)]E[E(X_j | \theta)] \\ &= E[\mu(\theta)^2] - E[\mu(\theta)]^2 = \text{Var}[\mu(\theta)] = a = 4/3. \end{aligned}$$

Question #237

Key: A

U	z	x	lognormal	with deductible
0.6217	0.31	5.8325	341.21	241.21
0.9941	2.52	7.49	1790.05	1690.05
0.8686	1.12	6.44	626.41	526.41
0.0485	-1.66	4.355	77.87	0
			Average	614.42

The value of z is obtained by inversion from the standard normal table. That is, $u = \Pr(Z \leq z)$. The value of x is obtained from $x = 0.75z + 5.6$. The lognormal value is obtained by exponentiating x and the final column applies the deductible.

Question #238

Key: B

$$\begin{aligned} \text{MSE} &= E[(X^2 - \theta^2)^2] = E(X^4 - 2X^2\theta^2 + \theta^4) \\ &= 24\theta^4 - 2(2\theta^2)\theta^2 + \theta^4 = 21\theta^4 \end{aligned}$$

Question #239

Key: C

The sample mean of $\frac{157(0) + 66(1) + 19(2) + 4(3) + 2(4)}{248} = 0.5$ is the maximum likelihood estimate of the geometric parameter β as well as the method of moments estimate of

the Poisson parameter λ . Then, $P = (1+0.5)^{-1} = 0.6667$ and $Q = e^{-0.5} = 0.6065$. The absolute difference is 0.0602.

Question #240

Key: D

$$\bar{x} = \frac{5000(0) + 2100(1) + 750(2) + 100(3) + 50(4)}{8000} = 0.5125 \text{ and}$$

$$s^2 = \frac{5000(0.5125)^2 + 2100(0.4875)^2 + 750(1.4875)^2 + 100(2.4875)^2 + 50(3.4875)^2}{7999} = 0.5874.$$

Then, $\hat{\mu} = \hat{v} = \bar{x} = 0.5125$ and $\hat{a} = s^2 - \bar{x} = 0.0749$. The credibility factor is

$$Z = \frac{1}{1 + 0.5125/0.0749} = 0.1275 \text{ and the estimate is } 0.1275(1) + 0.8725(0.5125) = 0.5747.$$

Question #241

Key: B

$s = F_n(3000) = 4/8 = 0.5$ because for the p - p plot the denominator is $n+1$.

$t = F(3000) = 1 - e^{-3000/3300} = 0.59711$. For the difference plot, D uses a denominator of n and so $D = 4/7 - 0.59711 = -0.02568$ and the answer is $0.5 - 0.59711 + 0.02568 = -0.071$.

Question #242**Key: B**

$\pi(q | 2, 2) \propto f(2 | q)f(2 | q)\pi(q) = q(q)(q^2 / 0.039) \propto q^4$. Because $\int_{0.2}^{0.5} q^4 dq = 0.006186$,

$\pi(q | 2, 2) = q^4 / 0.006186$. Given q , the expected number of claims is

$E(N | q) = 0(0.1) + 1(0.9 - q) + 2q = 0.9 + q$. The Bayesian estimate is

$$E(N | 2, 2) = \int_{0.2}^{0.5} (0.9 + q) \frac{q^4}{0.006186} dq = 1.319.$$

Question #243**Key: E**

$$0.689 = F_{500}(1500) = 0.5F_{500}(1000) + 0.5F_{500}(2000) = 0.5 \left(\frac{200 + 110}{500} + \frac{310 + x}{500} \right) \Rightarrow x = 69$$

$$0.839 = F_{500}(3500) = 0.5F_{500}(2000) + 0.5F_{500}(5000) = 0.5 \left(\frac{310 + 69}{500} + \frac{379 + y}{500} \right) \Rightarrow y = 81$$

Question #244**Key: A**

A is false because the test works best when the expected number of observations is about the same from interval to interval. B is true (*Loss Models*, 427-8), C is true (*Loss Models*, 428), and D is true (*Loss Models*, 430).

Question #245**Key: E**

$$n\lambda \geq \lambda_0 \left[1 + \left(\frac{\sigma_Y}{\theta_Y} \right)^2 \right]; \theta_Y = \alpha\theta = 10,000\alpha; \sigma_Y^2 = \alpha\theta^2 = 10^8\alpha$$

$$n\lambda \geq \left(\frac{1.96}{0.1} \right)^2 \left[1 + \frac{10^8\alpha}{10^8\alpha^2} \right] = 384.16(1 + \alpha^{-1})$$

Because α is needed, but not given, the answer cannot be determined from the information given.

Question #246**Key: E**

With $n + 1 = 16$, we need the $0.3(16) = 4.8$ and $0.65(16) = 10.4$ smallest observations.

They are $0.2(280) + 0.8(350) = 336$ and $0.6(450) + 0.4(490) = 466$.

The equations to solve are:

$$0.3 = 1 - \left(\frac{\theta^\gamma}{\theta^\gamma + 336^\gamma} \right)^2 \quad \text{and} \quad 0.65 = 1 - \left(\frac{\theta^\gamma}{\theta^\gamma + 466^\gamma} \right)^2$$

$$(0.7)^{-1/2} = 1 + (336/\theta)^\gamma \quad \text{and} \quad (0.35)^{-1/2} = 1 + (466/\theta)^\gamma$$

$$\frac{(0.7)^{-1/2} - 1}{(0.35)^{-1/2} - 1} = \frac{(336/\theta)^\gamma}{(466/\theta)^\gamma}$$

$$0.282814 = (336/466)^\gamma$$

$$\ln(0.282814) = \gamma \ln(336/466)$$

$$\gamma = 3.8614.$$

Question #247**Key: D**

Let E be the event of having 1 claim in the first four years. In four years, the total number of claims is Poisson(4λ).

$$\Pr(\text{Type I} | E) = \frac{\Pr(E | \text{Type I}) \Pr(\text{Type I})}{\Pr(E)} = \frac{e^{-1}(0.05)}{\Pr(E)} = \frac{0.01839}{\Pr(E)} = 0.14427$$

$$\Pr(\text{Type II} | E) = \frac{e^{-2}(2)(0.2)}{\Pr(E)} = \frac{0.05413}{\Pr(E)} = 0.42465$$

$$\Pr(\text{Type III} | E) = \frac{e^{-4}(4)(0.75)}{\Pr(E)} = \frac{0.05495}{\Pr(E)} = 0.43108$$

$$\text{Note: } \Pr(E) = 0.01839 + .05413 + .05495 = 0.12747$$

The Bayesian estimate of the number of claims in Year 5 is:

$$0.14427(0.25) + 0.42465(0.5) + 0.43108(1) = 0.67947.$$

Question #248**Key: B**

The sample mean is $0.2(400) + 0.7(800) + 0.1(1600) = 800$.

The sample variance is $0.2(400 - 800)^2 + 0.7(800 - 800)^2 + 0.1(1600 - 800)^2 = 96,000$.

The sample third central moment is

$$0.2(400 - 800)^3 + 0.7(800 - 800)^3 + 0.1(1600 - 800)^3 = 38,400,000.$$

The skewness coefficient is $38,400,000 / 96,000^{1.5} = 1.29$.

Question #249**Key: C**

Because $0.656 < 0.7654 < 0.773$, the simulated number of losses is 4. To simulate a loss by inversion, use

$$F(x) = 1 - e^{-(x/\theta)^\tau} = u$$

$$1 - u = e^{-(x/\theta)^\tau}$$

$$\ln(1 - u) = -(x/\theta)^\tau$$

$$x = \theta(-\ln(1 - u))^{1/\tau} = 200(-\ln(1 - u))^{1/2}$$

$$u_1 = 0.2738, x_1 = 113.12$$

$$u_2 = 0.5152, x_2 = 170.18$$

$$u_3 = 0.7537, x_3 = 236.75$$

$$u_4 = 0.6481, x_4 = 204.39$$

With a deductible of 150, the first loss produces no payments and 113.12 toward the 500 limit. The second loss produces a payment of 20.18 and the insured is now out-of-pocket 263.12. The third loss produces a payment of 86.75 and the insured is out 413.12. The deductible on the fourth loss is then 86.88 for a payment of $204.29 - 86.88 = 117.51$.

The total paid by the insurer is $20.18 + 86.75 + 117.51 = 224.44$.

Question #250**Key: A**

The density function is $f(x) = \theta x^{-2} e^{-\theta/x}$ and the likelihood function is

$$L(\theta) = \theta(186^{-2})e^{-\theta/186}\theta(91^{-2})e^{-\theta/91}\theta(66^{-2})e^{-\theta/66}(e^{-\theta/60})^7$$

$$\propto \theta^3 e^{-0.148184\theta}$$

$$l(\theta) = \ln L(\theta) = 3 \ln(\theta) - 0.148184\theta$$

$$l'(\theta) = 3\theta^{-1} - 0.148184 = 0$$

$$\theta = 3/0.148184 = 20.25.$$

The mode is $\theta/2 = 20.25/2 = 10.125$.

Question #251**Key: D**

We have $\mu(\theta) = 4\theta$ and $\mu = 4E(\theta) = 4(600) = 2400$. The average loss for Years 1 and 2 is 1650 and so $1800 = Z(1650) + (1-Z)(2400)$ which gives $Z = 0.8$. Because there were two years, $Z = 0.8 = 2/(2+k)$ which gives $k = 0.5$.

For three years, the revised value is $Z = 3/(3+0.5) = 6/7$ and the revised credibility estimate (using the new sample mean of 2021), $(6/7)(2021) + (1/7)(2400) = 2075.14$.

Question #252**Key: B**

The uncensored observations are 4 and 8 (values beyond 11 are not needed). The two r values are 10 and 5 and the two s values are 2 and 1. The Kaplan-Meier estimate is

$$\hat{S}(11) = (8/10)(4/5) = 0.64 \text{ and Greenwood's estimate is } (0.64)^2 \left(\frac{2}{10(8)} + \frac{1}{5(4)} \right) = 0.03072.$$

Question #253**Key: E**

$S_m | Q \sim \text{bin}(m, Q)$ and $Q \sim \text{beta}(1, 99)$. Then

$E(S_m) = E[E(S_m | Q)] = E(mQ) = m \frac{1}{1+99} = 0.01m$. For the mean to be at least 50, m must be at least 5,000.

Question #254**Key: D**

The posterior distribution is

$\pi(\lambda | \text{data}) \propto (e^{-\lambda})^{90} (\lambda e^{-\lambda})^7 (\lambda^2 e^{-\lambda})^2 (\lambda^3 e^{-\lambda}) \frac{\lambda^4 e^{-50\lambda}}{\lambda} = \lambda^{17} e^{-150\lambda}$ which is a gamma distribution

with parameters 18 and 1/150. For one risk, the estimated value is the mean, 18/150. For 100 risks it is 100(18)/150 = 12.

Alternatively,

The prior distribution is gamma with $\alpha = 4$ and $\beta = 50$. The posterior will be continue to be gamma, with $\alpha' = \alpha + \text{no. of claims} = 4 + 14 = 18$ and $\beta' = \beta + \text{no. of exposures} = 50 + 100 = 150$. Mean of the posterior = $\alpha / \beta = 18/150 = 0.12$. Expected number of claims for the portfolio = $0.12 (100) = 12$.

Question #255**Key: E**

$$0.95 = \Pr(0.95\mu < \bar{X} < 1.05\mu)$$

$$\bar{X} \sim N(\mu, \sigma^2/n = 1.44\mu^2/n)$$

$$0.95 = \Pr\left(\frac{0.95\mu - \mu}{1.2\mu/\sqrt{n}} < Z < \frac{1.05\mu - \mu}{1.2\mu/\sqrt{n}}\right)$$

$$0.95 = \Pr(-0.05\sqrt{n}/1.2 < Z < 0.05\sqrt{n}/1.2)$$

$$0.05\sqrt{n}/1.2 = 1.96$$

$$n = 2212.76.$$

Question #256**Key: B**

$$L(q) = \left[\binom{2}{0} (1-q)^2 \right]^{5000} \left[\binom{2}{1} q(1-q) \right]^{5000} = 2^{5000} q^{5000} (1-q)^{15000}$$

$$l(q) = 5000 \ln(2) + 5000 \ln(q) + 15000 \ln(1-q)$$

$$l'(q) = 5000q^{-1} - 15000(1-q)^{-1} = 0$$

$$\hat{q} = 0.25$$

$$l(0.25) = 5000 \ln(2) + 5000 \ln(0.25) + 15000 \ln(0.75) = -7780.97.$$

Question #257**Key: C**

The estimate of the overall mean, μ , is the sample mean, per vehicle, which is $7/10 = 0.7$.

With the Poisson assumption, this is also the estimate of $v = EPV$. The means for the two insureds are $2/5 = 0.4$ and $5/5 = 1.0$. The estimate of a is the usual non-parametric estimate,

$$VHM = \hat{a} = \frac{5(0.4 - 0.7)^2 + 5(1.0 - 0.7)^2 - (2-1)(0.7)}{10 - \frac{1}{10}(25 + 25)} = 0.04$$

(The above formula: Loss Models page 596, Herzog page 116, Dean page 25)

Then, $k = 0.7/0.04 = 17.5$ and so $Z = 5/(5+17.5) = 2/9$. The estimate for insured A is $(2/9)(0.4) + (7/9)(0.7) = 0.6333$.

Question #258**Key: A**

Item (i) indicates that X must one of the four given values.

Item (ii) indicates that X cannot be 200

Item (iii) indicates that X cannot be 400.

First assume $X = 100$. Then the values of r are 5, 3, 2, and 1 and the values of s are 2,

1, 1, and 1. Then $\hat{H}(410) = \frac{2}{5} + \frac{1}{3} + \frac{1}{2} + \frac{1}{1} = 2.23$ and thus the answer is 100. As a check,

if $X = 300$, the r values are 5, 4, 3, and 1 and the s values are 1, 1, 2, and 1. Then,

$$\hat{H}(410) = \frac{1}{5} + \frac{1}{4} + \frac{2}{3} + \frac{1}{1} = 2.12.$$

Question #259**Key: B**

The estimator of the Poisson parameter is the sample mean. Then,

$$E(\hat{\lambda}) = E(\bar{X}) = \lambda$$

$$\text{Var}(\hat{\lambda}) = \text{Var}(\bar{X}) = \lambda/n$$

$$c.v. = \sqrt{\lambda/n} / \lambda = 1/\sqrt{n\lambda}$$

It is estimated by $1/\sqrt{n\lambda} = 1/\sqrt{39} = 0.1601$.

Question #260**Key: E**

$$\begin{aligned} \Pr(\theta = 8 | X_1 = 5) &= \frac{\Pr(X_1 = 5 | \theta = 8) \Pr(\theta = 8)}{\Pr(X_1 = 5 | \theta = 8) \Pr(\theta = 8) + \Pr(X_1 = 5 | \theta = 2) \Pr(\theta = 2)} \\ &= \frac{0.125e^{-5(0.125)}(0.8)}{0.125e^{-5(0.125)}(0.8) + 0.5e^{-5(0.5)}(0.2)} = 0.867035. \end{aligned}$$

Then,

$$E(X_2 | X_1 = 5) = E(\theta | X_1 = 5) = 0.867035(8) + 0.132965(2) = 7.202.$$

Question #261**Key: D**

We have $q^{(T)} = 1 - (1 - q^{(1)})(1 - q^{(2)})$ and so $q^{(1)} = 1 - \frac{1 - q^{(T)}}{1 - q^{(2)}} = 1 - \frac{1 - q^{(T)}}{1 - 0.05} = \frac{q^{(T)} - 0.05}{0.95}$.

Then, ${}_{20}q_0^{(1)} = 0.05/0.95 = 0.05263$, ${}_{20}q_{20}^{(1)} = 0.132/0.95 = 0.1389$, and

${}_{40}p_0^{(1)} = 0.9474(0.8611) = 0.8158$. Out of 1000 at age 0, 816 are expected to survive to age 40.

Question #262**Key: D**

$$L(\omega) = \frac{\frac{1}{\omega} \frac{1}{\omega} \frac{1}{\omega} \left(\frac{\omega-4-p}{\omega} \right)^2}{\left(\frac{\omega-4}{\omega} \right)^5} = \frac{(\omega-4-p)^2}{(\omega-4)^5}$$

$$l(\omega) = 2 \ln(\omega-4-p) - 5 \ln(\omega-4)$$

$$l'(\omega) = \frac{2}{\omega-4-p} - \frac{5}{\omega-4} = 0$$

$$0 = l'(29) = \frac{2}{25-p} - \frac{5}{25}$$

$$p = 15.$$

The denominator in the likelihood function is $S(4)$ to the power of five to reflect the fact that it is known that each observation is greater than 4.

Question #263**Key: B**

$$\mu(\lambda) = v(\lambda) = \lambda$$

$$\mu = v = E(\lambda) = 0.1\Gamma(1+1/2) = 0.088623$$

$$\text{VHM} = a = \text{Var}(\lambda) = (0.1)^2 \Gamma(1+2/2) - 0.088623^2 = 0.002146$$

$$Z = \frac{500}{500 + 0.088623/0.002146} = 0.92371.$$

The estimate for one insured for one month is

$$0.92371(35/500) + 0.07629(0.088623) = 0.07142. \quad \text{For 300 insureds for 12 months it is } (300)(12)(0.07142) = 257.11.$$

Question #264**Key: D**

With no censoring the r values are 12, 9, 8, 7, 6, 4, and 3 and the s values are 3, 1, 1, 1, 2, 1, 1 (the two values at 7500 are not needed). Then,

$$\hat{H}_1(7000) = \frac{3}{12} + \frac{1}{9} + \frac{1}{8} + \frac{1}{7} + \frac{2}{6} + \frac{1}{4} + \frac{1}{3} = 1.5456.$$

With censoring, there are only five uncensored values with r values of 9, 8, 7, 4, and 3 and all five s values are 1. Then,

$$\hat{H}_2(7000) = \frac{1}{9} + \frac{1}{8} + \frac{1}{7} + \frac{1}{4} + \frac{1}{3} = 0.9623. \quad \text{The absolute difference is 0.5833.}$$

Question #265**Key: A**

The simulated paid loss is $\exp[0.494\Phi^{-1}(u)+13.294]$ where $\Phi^{-1}(u)$ is the inverse of the standard normal distribution function. The four simulated paid losses are 450,161, 330,041, 939,798, and 688,451 for an average of 602,113. The multiplier for unpaid losses is $0.801(2006-2005)^{0.851}e^{-0.747(2006-2005)} = 0.3795$ and the answer is $0.3795(602,113) = 228,502$

Question #266**Key: A**

The deduction to get the SBC is $(r/2)\ln(n) = (r/2)\ln(260) = 2.78r$ where r is the number of parameters. The SBC values are then -416.78, -417.56, -419.34, -420.12, and -425.68. The largest value is the first one, so model I is to be selected.

Question #267**Key: E**

$$\begin{aligned}\Pr(\lambda = 1 | X_1 = r) &= \frac{\Pr(X_1 = r | \lambda = 1)\Pr(\lambda = 1)}{\Pr(X_1 = r | \lambda = 1)\Pr(\lambda = 1) + \Pr(X_1 = r | \lambda = 3)\Pr(\lambda = 3)} \\ &= \frac{\frac{e^{-1}}{r!}(0.75)}{\frac{e^{-1}}{r!}(0.75) + \frac{e^{-3}3^r}{r!}(0.25)} = \frac{0.2759}{0.2759 + 0.1245(3^r)}.\end{aligned}$$

Then,

$$\begin{aligned}2.98 &= \frac{0.2759}{0.2759 + 0.1245(3^r)}(1) + \frac{0.1245(3^r)}{0.2759 + 0.1245(3^r)}(3) \\ &= \frac{0.2759 + 0.3735(3^r)}{0.2759 + 0.1245(3^r)}.\end{aligned}$$

Rearrange to obtain

$$0.82218 + 0.037103(3^r) = 0.2759 + 0.03735(3^r)$$

$$0.54628 = 0.00025(3^r)$$

$$r = 7.$$

Because the risks are Poisson, ($\mu = \text{EPV}$, $a = \text{VHM}$):

$$\mu = v = E(\lambda) = 0.75(1) + 0.25(3) = 1.5$$

$$a = \text{Var}(\lambda) = 0.75(1) + 0.25(9) - 2.25 = 0.75$$

$$Z = \frac{1}{1 + 1.5/0.75} = 1/3$$

and the estimate is $(1/3)(7) + (2/3)(1.5) = 3.33$.

Question #268**Key: E**

The uniform kernel spreads the probability of 0.1 to 10 units each side of an observation. So the observation at 25 contributes a density of 0.005 from 15 to 35, contributing nothing to survival past age 40. The same applies to the point at 30. For the next 7 points:

35 contributes probability from 25 to 45 for $5(0.005) = 0.025$ above age 40.

35 contributes probability from 25 to 45 for $5(0.005) = 0.025$ above age 40.

37 contributes probability from 27 to 47 for $7(0.005) = 0.035$ above age 40.

39 contributes probability from 29 to 49 for $9(0.005) = 0.045$ above age 40.

45 contributes probability from 35 to 55 for $15(0.005) = 0.075$ above age 40.

47 contributes probability from 37 to 57 for $17(0.005) = 0.085$ above age 40.

49 contributes probability from 39 to 59 for $19(0.005) = 0.095$ above age 40.

The observation at 55 contributes all 0.1 of probability. The total is 0.485.

Question #269**Key: A**

$$X \sim \text{Exp}(\theta)$$

$$\sum_{i=1}^n X_i \sim \Gamma(n, \theta)$$

$$\bar{X} \sim \Gamma(n, \theta/n)$$

$$E(\bar{X}^2) = (\theta/n)^2(n)(n+1) = (n+1)\theta^2/n.$$

The second line follows because an exponential distribution is a gamma distribution with $\alpha = 1$ and the sum of independent gamma random variables is gamma with the “ α ” parameters added. The third line follows because the gamma distribution is a scale distribution. Multiplying by $1/n$ retains the gamma distribution with the “ θ ” parameter multiplied by $1/n$.

Question #270**Key: C**

The sample means are 3, 5, and 4 and the overall mean is 4. Then,

$$\hat{v} = \frac{1+0+0+1+0+0+1+1+1+1+1+1}{3(4-1)} = \frac{8}{9}$$

$$\hat{a} = \frac{(3-4)^2 + (5-4)^2 + (4-4)^2}{3-1} - \frac{8/9}{4} = \frac{7}{9} = 0.78.$$

Question #271
DELETED

Question #272
Key: C

$$\pi(q | 2) = 6q^2(1-q)^2 6q(1-q) \propto q^3(1-q)^3$$

The mode can be determined by setting the derivative equal to zero.

$$\pi'(q | 2) \propto 3q^2(1-q)^3 - 3q^3(1-q)^2 = 0$$

$$(1-q) - q = 0$$

$$q = 0.5.$$

Question #273
Key: B

For the severity distribution the mean is 5,000 and the variance is $10,000^2/12$. For credibility based on accuracy with regard to the number of claims,

$$2000 = \left(\frac{z}{0.03} \right)^2, \quad z^2 = 1.8$$

where z is the appropriate value from the standard normal distribution. For credibility based on accuracy with regard to the total cost of claims, the number of claims needed is

$$\frac{z^2}{0.05^2} \left(1 + \frac{10000^2/12}{5000^2} \right) = 960.$$

Question #274**Key: C**

$$\hat{S}(10) = e^{-\hat{H}(10)} = 0.575$$

$$\hat{H}(10) = -\ln(0.575) = 0.5534 = \frac{1}{50} + \frac{3}{49} + \frac{5}{k} + \frac{7}{12}.$$

The solution is $k = 36$.

Question #275**Key: A**

The annual dental charges are simulated from

$$u = 1 - e^{-x/1000}$$

$$x = -1000 \ln(1 - u).$$

The four simulated values are 356.67, 2525.73, 1203.97, and 83.38. The reimbursements are 205.34 (80% of 256.67), 1000 (the maximum), 883.18 (80% of 1103.97), and 0. The total is 2088.52 and the average is 522.13.

Question #276**Key: B**

$$L(\theta) = \left(1 - \frac{\theta}{10}\right)^9 \left(\frac{\theta}{10} - \frac{\theta}{25}\right)^6 \left(\frac{\theta}{25}\right)^5 \propto (10 - \theta)^9 \theta^{11}$$

$$l(\theta) = 9 \ln(10 - \theta) + 11 \ln(\theta)$$

$$l'(\theta) = -\frac{9}{10 - \theta} + \frac{11}{\theta} = 0$$

$$11(10 - \theta) = 9\theta$$

$$110 = 20\theta$$

$$\theta = 110/20 = 5.5.$$

Question #277**Key: A**

The maximum likelihood estimate is $\hat{\theta} = \bar{x} = 1000$. The quantity to be estimated is $S(\theta) = \exp(-1500/\theta)$ and $S'(\theta) = 1500\theta^{-2} \exp(-1500/\theta)$. For the delta method,

$$\begin{aligned} \text{Var}[S(\hat{\theta})] &\cong [S'(\hat{\theta})]^2 \text{Var}(\hat{\theta}) \\ &= [1500(1000)^{-2} \exp(-1500/1000)]^2 (1000^2 / 6) \\ &= 0.01867. \end{aligned}$$

This is based on $\text{Var}(\hat{\theta}) = \text{Var}(\bar{X}) = \text{Var}(X) / n = \theta^2 / n$.

Question #278**Key: A**

Based on the information given

$$0.21 = \frac{36}{n} + \frac{0.4x}{n}$$

$$0.51 = \frac{36}{n} + \frac{x}{n} + \frac{0.6y}{n}$$

$$n = 200 + x + y.$$

Then,

$$0.21(200 + x + y) = 36 + 0.4x$$

$$0.51(200 + x + y) = 36 + x + 0.6y$$

and these linear equations can be solved for $x = 119.37$.

Question #279**Key: B**

Pays 80% of loss over 20, with cap of payment at 60, hence cap of loss of

$$\frac{60}{0.8} + 20 = 95 = u$$

$$\text{Th 5.13 } E(Y \text{ per loss}) = \alpha [E(X \wedge 95) - E(X \wedge 20)]$$

$$= 0.8 \left[\int_0^{95} S(x) dx - \int_0^{20} S(x) dx \right]$$

$$= 0.8 \int_{20}^{95} S(x) dx = 0.8 \int_{20}^{95} \left(1 - \frac{x^2}{10,000} \right) dx = 0.8 \left(x - \frac{x^3}{30,000} \Big|_{20}^{95} \right) = (0.8)(46.6875) = 37.35$$

$$E(Y \text{ per payment}) = \frac{E(Y \text{ per loss})}{1 - F(20)} = \frac{37.35}{0.96} = 38.91$$

Question #280**Key: D**Let S = aggregate claims, I_5 = claims covered by stop loss

S	I_5	
0	0	
5	0	
x	x - 5	(i.e., for any $S > 5$, $I_5 = S - 5$)

$$E[I_5] = E[S] - 5 + 5 \Pr(0 \text{ claims})$$

$$E[S] = 5 \times (0.6 \times 5 + 0.4 \times k) = 15 + 2k$$

$$\Pr(0 \text{ claims}) = e^{-5}$$

$$E[I_5] = 15 + 2k - 5 + 5e^{-5} = 28.03$$

$$10.034 + 2k = 28.03$$

$$2k = 18$$

$$k = 9$$

Question #281**Key: B**Let R = Equity index return. $R \sim N(\mu = 8\%, \sigma = 16\%)$ $X = 0.75 R$ $X \sim N(\mu = 6\%, \sigma = 12\%)$ Y = crediting rate = $\text{Max}(R, 3\%)$

$$Y = X + 3\% - (X \wedge 3\%)$$

(If $X < 3\%$, $X \wedge 3\% = X$ and $Y = X + 3\% - X = 3\%$)(If $X > 3\%$ $X \wedge 3\% = 3\%$ and $Y = X + 3\% - 3\% = X$)

$$\begin{aligned} E(Y) &= E(X + 3\% - (X \wedge 3\%)) \\ &= E(X) + 3\% - E(X \wedge 3\%) \\ &= 6\% + 3\% - (-0.43\%) \\ &= 9.43\% \end{aligned}$$

Note that $E(X \wedge 3\%)$ is a table lookup in the given information.

Question #282**Key: A**Let S = aggregate losses, X = severity

Since the frequency is Poisson,

$$\text{Var}(S) = \lambda E(X^2)$$

$$E(X^2) = \frac{2^2 \Gamma(3) \Gamma(1)}{\Gamma(3)} = 4 \quad (\text{table lookup})$$

$$\text{Var}(S) = 3 \times 4 = 12$$

You would get the same result if you used

$$\text{Var}(S) = E(N)\text{Var}(X) + \text{Var}(N)(E(X))^2$$

Question #283**Key: D**For each member $P(z) = [1 - 1.5(z-1)]^{-1}$ so for family of 4 $P(z) = [1 - 1.5(z-1)]^{-4}$ negative binomial with $\beta = 1.5$ $r = 4$

<u>k</u>	<u>P_k</u>
0	0.026
1	0.061
2	0.092
3+	0.821

$$E(N \wedge 3) = 0 \times 0.026 + 1 \times 0.061 + 2 \times 0.092 + 3 \times 0.821 = 2.71$$

$$E(N) - E(N \wedge 3) = 6 - 2.71 = 3.29$$

$$3.29 \times 100 \text{ per visit} = 329$$

Alternatively, without using probability generating functions, a geometric distribution is a special case of the negative binomial with $r = 1$.Summing four independent negative binomial distributions, each with $\beta = 1.5$ and $r = 1$ gives a negative binomial distribution with $\beta = 1.5$ and $r = 4$. Then continue as above.

Question #284**Key: E**

$$\begin{aligned}
E(X \wedge 2) &= 1 \times f(1) + 2(1 - F(1)) \\
&= 1 \times f(1) + 2(1 - f(0) - f(1)) \\
&= 1 \times 3e^{-3} + 2(1 - e^{-3} - 3e^{-3}) \\
&= 2 - 5e^{-3} \\
&= 1.75
\end{aligned}$$

$$\begin{aligned}
\text{Cost per loss with deductible} &= E(X) - E(X \wedge 2) \\
&= 3 - 1.75 \\
&= 1.25
\end{aligned}$$

$$\text{Cost per loss with coinsurance} = \alpha E(X) = 3\alpha$$

$$\begin{aligned}
\text{Equating cost, } 3\alpha &= 1.25 \\
\alpha &= 0.42
\end{aligned}$$

Question #285**Key: A**

Let N be the number of clubs accepted
 X be the number of members of a selected club
 S be the total persons appearing

N is binomial with $m = 1000$ $q = 0.20$

$$E(N) = (1000)(0.20) = 200$$

$$\text{Var}(N) = (1000)(0.20)(0.80) = 160$$

$$E(S) = E(N)E(X) = (200)(20) = 4000$$

$$\begin{aligned}
\text{Var}(S) &= E(N) \text{Var}(X) + \text{Var}(N)[E(X)]^2 \\
&= (200)(20) + (160)(20)^2 \\
&= 68,000
\end{aligned}$$

$$\begin{aligned}
\text{Budget} &= 10 \times E(S) + 10 \times \sqrt{\text{Var}(S)} \\
&= 10 \times 4000 + 10 \times \sqrt{68,000} \\
&= 42,610
\end{aligned}$$

Question #286

Key: C

Insurance pays 80% of the portion of annual claim between 6,000 and 1,000, and 90% of the portion of annual claims over 14,000.

The 14,000 breakpoint is where Michael reaches 10,000 that he has paid:

1000 = deductible

1000 = 20% of costs between 1000 and 6000

8000 = 100% of costs between 14,000 and 6,000

$$E[X \wedge x] = \theta \left(1 - \frac{\theta}{x + \theta} \right) = \frac{5000x}{x + 5000}$$

x	$E[X \wedge x]$
1000	833.33
6000	2727.27
14000	3684.21
∞	5000

$$0.80[E[X \wedge 6000] - E[X \wedge 1000]] + 0.90[E[X] - E[X \wedge 14000]]$$

$$\Rightarrow 0.80[2727.27 - 833.33] + 0.90[5000 - 3684.21]$$

$$= 1515.15 + 1184.21 = 2699.36$$

$$\Rightarrow 2700$$

Question #287

Key: D

We have the following table:

Item	Dist	$E()$	$Var()$
Number claims	$NB(16, 6)$	$16 \times 6 = 96$	$16 \times 6 \times 7 = 672$
Claims amounts	$U(0, 8)$	$8 / 2 = 4$	$8^2 / 12 = 5.33$
Aggregate		$4 \times 96 = 384$	$96 \times 5.33 + 672 \times 4^2 = 11,264$

$$\text{Premium} = E(S) + 1.645 * \text{Sqrt}(Var(S)) =$$

$$= 384 + 1.645 * \text{Sqrt}(11,264)$$

$$= 559$$

1.645 is the 95th percentile of the standard normal distribution.

Question #288**Key: E**

With probability p , $\text{Prob}(N = 2) = 0.5^2 = 0.25$

With probability $(1 - p)$, $\text{Prob}(N = 2) = \text{Combin}(4, 2) * 0.5^4 = 0.375$

$$\text{Prob}(N = 2) = p \times 0.25 + (1 - p) \times 0.375$$

$$0.375 - 0.125 p$$

Question #289**Key: D**

600 can be obtained only 2 ways, from $500 + 100$ or from 6×100 .

Since $\lambda = 5$ and $p(100) = 0.8$, $p(500) = 0.16$,

$$p(6 \text{ claims for } 100) = \frac{e^{-5} 5^6}{6!} (0.8)^6 = 0.03833 \text{ or } 3.83\%$$

$$p(500 + 100) = \frac{e^{-5} 5^2}{2!} \left[(0.8)^1 (0.16)^1 (2) \right] = 0.02156 = 2.16\%$$

The factor of 2 inside the bracket is because you could get a 500 then 100 or you could get a 100 then 500.

$$\text{Total} = 3.83\% + 2.16\% = 5.99\%$$